

## Modulation of human ovarian function by melatonin

Seema Rai<sup>1</sup>, Hindole Gosh<sup>2</sup>

<sup>1</sup>Department of Zoology, Guru Ghasidas Vishwavidyalaya, Bilaspur, Chhattisgarh, India, <sup>2</sup>Department of Zoology, Kalinga University, Raipur, Chhattisgarh, India

### TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Regulation of pineal melatonin biosynthesis
4. Influence of rhythmicity of melatonin during onset of puberty
5. Modulation of ovarian function by melatonin
  - 5.1. Distribution of melatonin receptors in the ovary
  - 5.2. Modulation of follicular growth by melatonin
  - 5.3. Influence of melatonin on ovarian steroidogenesis
  - 5.4. Effect of melatonin on ovulation
  - 5.5. Effect of melatonin in maintenance of oocyte quality
6. Modulation of luteal function by melatonin
7. Protective effect of melatonin in embryo culture
8. Effect of melatonin in success rate of IVF
9. Conclusion
10. Acknowledgments
11. References

### 1. ABSTRACT

Melatonin, a hormone which is primarily released by the pineal gland, has a wide range of actions in the female reproductive tract. While the melatonin receptor subtype, MT<sub>3</sub>, has been identified in amphibian animals and birds, in humans and other mammals, melatonin acts through, MT<sub>1</sub> and MT<sub>2</sub> receptor subtypes which are expressed in human ovaries. The rhythmic release of melatonin starts at puberty and continues throughout fertile female life, affecting and regulating diverse ovarian functions. Here, we discuss the importance of melatonin in regulating folliculogenesis, oocyte quality, ovulation and luteal function, sex steroid receptor gene expression, ovarian steroidogenesis including the production and steroidogenic enzyme activities in the egg and thecal cells. Melatonin improves the egg quality and increases the chance of success of *in vitro* fertilization (IVF). In view of such extensive actions, melatonin is central to the fertility in females.

The objective of this review is to recapitulate the current understanding of the role of melatonin and its receptors.

### 2. INTRODUCTION

Melatonin (N-acetyl-5-methoxytryptamine), was identified 60 years ago as a neurohormone chiefly secreted by the pineal gland (epiphysis cerebri). Since then, various pinealologists have revealed the physiology and biochemistry of the pineal melatonin. Reports indicate, that its interaction is not limited to the endocrine organs, rather melatonin interacts with other non-endocrine tissues and hence participates in the regulation of various metabolic, immunologic, reproductive, and, other vital physiologic processes coordinating with neuroendocrine network system (1-4). Melatonin is an indoleamine, a tryptophan derivative, which is an

essential amino acid, synthesized via serotonin and acting as a neurohormone.


The duration of melatonin secretion tracks the dark phase of light /dark cycle but, in the case of the mammals this humoral signal is considered a crucial component of the mechanism governing mammalian photoperiodism. After its release in the circulation, it acts as a chemical messenger of photoperiodic phases of the light / dark cycle and regulates the circadian rhythm, sleep-wake cycle as well as various endocrine processes or factors. Melatonin due to its rhythmic diurnal and/or nocturnal fluctuations is assumed to correspond circadian activities with different photoperiods and hormones of the hypothalamic-pituitary axis (5-7). Moreover, the changes ensure the duration and the magnitude i.e. diurnal (clock) and seasonal (calendar) of circulatory levels of the hormone in the blood. Melatonin throughout the year provides information about the season and hence, it is popularly considered as a clock and calendar. This hormone contributes to various central and peripheral physiological actions related to circadian rhythm and reproduction. It participates in the regulation of reproduction in seasonal breeders and phasing of circadian rhythm in mammalian species (8-10). Pieces of evidence categorize it as an immune-regulatory, antioxidative, anti-stress, antidiabetic, neuroprotective, hepatic, and nephroprotective hormone (11-16). By the virtue of being a highly conserved and dispersed biologically active small lipophilic indoleamine, melatonin can act either via receptor (G-protein coupled melatonin receptors) or non- receptor-mediated pathway (17).

Even though this neurohormone is chiefly synthesized and secreted in the circulation by the pineal gland. There are reports providing evidence of its synthesis by other organs and tissues like the retina, the hardierian gland, lymphocytes, gastrointestinal tract, and skin, which is termed as extra pineal melatonin (3, 17 -19). Available literature suggests that melatonin may regulate the reproductive neuroendocrine axis in humans (20). The goal of the present review is to explore the comprehensive information about the new horizon of melatonin research, pointing specifically about its modulatory function in female reproductive

physiology, with the hypothesis of a molecular mechanistic approach to stimulate future research. It is expected that this may throw light on therapeutic potentials of melatonin which would be beneficial in medical science for the improvement of human reproductive health and fertility.

### 3. REGULATION OF PINEAL MELATONIN BIOSYNTHESIS

Melatonin is a small (molecular weight 232.3) indoleamine, secreted rhythmically being with the highest peak during the dark period of light/dark cycle. Alterations in environmental lighting are signaled as multisynaptic neural inputs by the central nervous system *via* peripheral nerves into a hormonal output of the pineal gland. In mammals, circadian photoreceptors of the retina convert light and darkness into signals that are sent directly to the SCN through the main pathway, the Retino Hypothalamic Tract (RHT). From the SCN, neuronal projections make synaptic connections in the PVN of the hypothalamus descending onward through the medial forebrain bundle to the intermediolateral cell column of the spinal cord from where preganglionic fibers reach the superior cervical ganglia. Sympathetic postganglionic noradrenergic fibers, from the superior cervical ganglia, innervate the pineal gland through the *nervi conarii*. Interruption of this regulation pathway by a lesion of the SCN or PVN, or by superior cervical ganglionectomy abolishes the pineal gland synthesis. Thus, the production of pineal melatonin occurs in response to nor-adrenergic stimulation which produces a cascade of biochemical events within the pinealocytes. The *N*-acetyltransferase (NAT) activity represents a key regulatory step in melatonin synthesis. Noradrenaline release, from the sympathetic nerves that innervate the pineal gland, is normally high at night and low during the day. In most species, noradrenaline interacts with both beta 1- and alpha1-adrenergic receptors present in the pineal gland. In the rat pinealocyte, stimulation of adrenergic receptors induces a rise in adenylate cyclase, and the cyclic adenosine 3',5'-monophosphate (cAMP) signaling pathway activates the NAT enzyme that catalyzes the rate-limiting step of melatonin synthesis. Simultaneous activation of alpha1-receptors potentiates the effect




**Figure 1.** Pathway showing biosynthesis of melatonin and its regulation. Figure 2. Nor-epinephrine (NE) from sympathetic nerve fibre is the key regulator for melatonin synthesis and the key regulator of production of melatonin in the dark and light phases. Gamma-aminobutyric acid (GABA) released by neurons of SCN (suprachiasmatic nuclei) into the paraventricular nucleus (PVN) and switch off the release of melatonin from pineal gland during day time (Upper Panel). In night time NE released from superior cervical ganglia (SCG) activating the projection of PVN to the pre-ganglionic cell into the intermediolateral cell column (ILC) of the spinal cord. This further stimulates the  $\beta_1$ -adrenergic receptors present in the pinealocytes and promotes protein kinase A (PKA) stimulation, consequently, escalation in the AANAT enzyme activity leads to synthesis and release of melatonin in circulation (Lower panel).

mediated through the beta1-receptors (21). Hence, the rhythmic melatonin synthesis is ensured by the oscillating cAMP-dependent transcriptional control mechanism. Melatonin's biosynthetic pathway involves tryptophan, which is hydroxylated and decarboxylated to serotonin, and then serotonin is acetylated by the rate-limiting enzyme NAT and further methylated by hydroxy indole-O-methyltransferase to melatonin (5-methoxy-N-acetyltryptamine) in the pineal gland. After the synthesis, melatonin is secreted into the blood and cerebrospinal fluid (Figure 1).

## 4. INFLUENCE OF RHYTHMICITY OF MELATONIN DURING ONSET OF PUBERTY

There is controversy regarding the effect of melatonin during the pre and post-pubertal stage. As reviewed by Waldhauser and his associates in 1993,

before the age of 3 months, the circulatory concentration of melatonin remains either non-detectable or very low. It is only thereafter that, this hormone begins to synthesize (22). During the fetal and the first year of life of all individuals HPG axis remains active; afterward, it becomes quiescent till the age of 10 years. As long as the age of individual advances, the circulatory melatonin gradually drops down to a significantly decreased level and stimulates the enhancement of pulsatile secretion of the gonadotropin-releasing hormone as well as the reproductive axis resulting in the onset of puberty (23). Further, nuclear magnetic resonance (NMR) results also provide shreds of evidence of the independence of the production rate of melatonin with age and growth in pineal size between 1 to 15 years of age. Hence, decrement in the circulatory melatonin concentration provides a link between the increase in body mass / and / or due to sexual maturation (24). The hypothesis, regarding the correlation amid endogenous melatonin and puberty, based on the earlier finding suggests that prepubertal high circulatory level of melatonin results in inhibition of secretion of gonadotropin hormones (25). Afterward, as the individual progresses to the age of puberty, endogenous melatonin comes down to that critical level, which ultimately triggers or switches on the puberty. This hypothesis recommends that the melatonin influences not only the pubertal sexual maturation but even the gonadal and genital development of offspring in rats. However, such a hypothesis still seeks attention for more investigation to be evaluated further for humans (25). It may, therefore, be ventured that before puberty, melatonin concentrations remain too elevated consequently inhibiting activation of the hypothalamic axis. The important aspect to understand is just before puberty, circulatory melatonin drops even lower than the threshold value hence forming the trigger signals of GnRH from the hypothalamus which results in the onset of pubertal changes (25). In various studies, it has been shown that there is a negative correlation between the melatonin and LH concentration and also that it controls the pulsatile secretion of LH (25). Studies have demonstrated that high nocturnal melatonin secretion in children delays puberty whereas low levels of melatonin are associated with precocious puberty (26). Therefore, it is the decline of melatonin which causes the onset of puberty.



**Figure 2.** Diagrammatic representation of the proposed mechanism of action of melatonin following receptor and non-receptor mediated pathway. Melatonin in circulation taken up by TC, GC, enters to follicular fluid stimulated the LH, FSH mRNA expression and maturation inducing hormone activity via interacting with its own receptors (MT1R and MT2R) which in turn enhancing and regulating steroidogenic enzyme activities. ROS generation during follicular rupture results in ovulation and steroidogenesis (monooxygenase reaction for synthesis of Progesterone) in corpus luteum in because of inflammatory reaction. Circulatory melatonin released from pineal gland, acts as free radical scavenger and maintaining balance between ROS generation and antioxidative enzymes hence promoting mature ovum

## 5. MODULATION OF OVARIAN FUNCTION BY MELATONIN

A high concentration of melatonin in follicular fluid (FF), relative to the circulatory plasma level and expression of its both the receptors MT1 and MT2 in granulosa cells, luteal cells, antral follicles and corpus luteum of rats, indicates that it plays many essential roles in the regulation of mammalian reproductive processes (27-31). Further, literature provides evidence that melatonin alters granulosa cell steroidogenesis and follicular function in avians, rodents, and humans (32-35) (Figure 2).

### 5.1. Distribution of melatonin receptors in the ovary

The available report strongly suggests that neurohormone melatonin is neither

antigonadotrophic nor progonadotrophic. Therefore, it is very clear that change in nocturnal circulatory melatonin act as a messenger for passive signal updating the hypothalamic-pituitary-gonadal (HPG)-axis regarding the information about the time of year (36-37). Pineal melatonin is also documented for its participation in the regulation of reproduction in polyestrous mammals such as laboratory rats and human females. This documentation has been supported by the results of the decrement on gonadotrophins and impairments in the periodicity of estrous cycles (38-39). The melatonin actions are mediated through specific membrane receptors which are either G-protein-coupled receptors (GPCRs) or on nuclear binding sites; the latter corresponds to orphan members of the nuclear receptor RZR/ROR superfamily (40). Among the proposed three subtypes of mammalian membrane melatonin receptors, MT1 and MT2, are members of



the 7-transmembrane G protein-coupled receptor family (41-43). MT<sub>3</sub>, the third receptor which has all the features of a melatonin receptor, is an enzyme called quinone reductase 2, which has been identified only in a few animals (41-42). The signal transduction cascade associated with the activation of MT<sub>1</sub> or MT<sub>2</sub> in target cells results in the inhibition of adenylate cyclase activity (42). Activation of these receptors results in the reduction of forskolin-induced cyclic adenosine 3050 monophosphate (cAMP) formation leading to the subsequent reduction in activated protein kinase A (44). MT<sub>1</sub> and MT<sub>2</sub> receptors trigger the physiological activity following the conventional rule of biochemical pathways of signal transduction. Melatonin may activate different second messenger cascades by interacting with the same receptor subtype depending on the tissue, organ, and species. The literature report, therefore, demonstrates that melatonin may directly affect ovarian function as its receptor binding sites (MT<sub>1</sub> and MT<sub>2</sub> receptors) are present human GCs as well as on luteal cells (36-37).

### 5.2. Modulation of follicular growth by melatonin

Folliculogenesis is a complex process involving endocrine, paracrine, and autocrine mechanisms. It begins with the formation of numerous primordial follicles. Primordial follicles progressively develop into the primary, preantral, and antral stages, ultimately reach the pre-ovulatory stage and are capable of releasing an oocyte for subsequent fertilization. Depending upon the species, the process of folliculogenesis is critically dependent on circulating levels of FSH. Interestingly, of the vast number of follicles that are recruited from the ovarian follicular reserve to develop during each reproductive cycle, few (arguably <0.1%) of these follicles will ever fully mature and shed their ovum. Instead, all but the so-called selected follicles die due to atresia. The antral follicle is a small fluid-filled space, which culminates in the fully mature follicle. Human preovulatory follicular fluid FF has been reported with an almost threefold higher circulatory level of melatonin than normal (26-27, 34). Further, the presence of melatonin, its precursors, serotonin, and N-acetylserotonin, have also been documented in extracts of human ovary. Enzyme activities of NAT

and HIOMT required for melatonin synthesis, are also reported in human ovarian homogenates (45). These findings support the possible root cause of the presence of a higher concentration of melatonin in ovarian follicular fluid, suggesting that after its synthesis by the ovary, melatonin might have released into the FF. Nevertheless, the current investigations narrate the opinion that the majority of the detected ovarian and preovulatory FF melatonin is derived from the endogenous circulation (46). The scenario becomes a bit more apparent and it can be explained that the follicular melatonin is taken up from the circulation following its synthesis from the pineal gland. Reportedly, the process of folliculogenesis produces a lot of ROS because of ongoing functional and morphological changes in the two major gears ovarian follicle viz the theca and granulosa cells. Since melatonin is present in the vicinity of the ovarian follicle, therefore, and being a strong antioxidant, it protects the female gamete from the oxidative mutilation and hence regulates healthy folliculogenesis.

### 5.3. Influence of melatonin on ovarian steroidogenesis

The two-cell theory explains a tuned model of co-operativity of theca and granulosa cells in the process of ovarian steroid hormone synthesis, important for the estradiol production. Recruitment of antral follicles is characterized by the induction of expression of mRNAs encoding a range of steroidogenic enzymes, gonadotropin receptors, and local regulatory factors. The mechanism of selection of the graafian follicle seems to be linked to the timing of mRNA expression encoding LH receptors in GCs (47). *In vitro* melatonin supplementation (10 pM–100 nM) resulted in a remarkable increase in mRNA expression of LH (but not FSH) receptors in human GCs (31). When theca cells and GCs were separated, melatonin supplementation resulted in the decrement of the production of P by theca, irrespective of its influence on the GCs. It may be assumed that melatonin might be involved in suppressing follicular (thecal) steroidogenesis via modulating the cAMP level. Further, reports demonstrate that melatonin blocks the expression of the steroidogenic acute regulatory protein (48). It is believed that steroidogenic acute regulatory (StAR) protein

determines the translocation of cholesterol across the intermembrane space into the inner membrane where P450<sub>scc</sub> cleaves cholesterol into pregnenolone. Melatonin (10 nM) treatment for 3 hours reduced the steroidogenic acute regulatory protein expression stimulated by hCG in mouse Leydig tumor cells (48). Further, pinealectomy and/or administration of melatonin influences CYP17 expression and therefore the process of steroidogenesis as well (38). It seems that there is a complex effect of melatonin on follicular steroid production depending on the cell type (theca cell or GC), duration of treatment (acute or long-term response), experimental model (cell culture or follicle culture), species, and dose (48).

Melatonin treatment with a dosage of 0.01–10 mg/mL stimulates IGF-I production, reported by cultured human GCs (49). Picinato and his associates during 2008 demonstrated that melatonin (0.1 mM) induces the IGF-I receptor and activates two intracellular signaling pathways, the PI3 K/AKT, which is mainly involved with cell metabolism, and MEK/ERKs that participate in cell proliferation, growth, and differentiation (50). Insulin-like growth factors are produced by GCs and theca cells during follicle development (51). Insulin-like growth factors are mitogenic and antiapoptotic peptides that promote differentiation and also possess insulin-like metabolic effects mediated by binding to specific high-affinity membrane receptors. The IGF-I and IGF-II stimulate DNA synthesis and E2 and P secretion by human GCs and granulosa-luteal cells (51). Insulin-like growth factor-I is antiapoptotic in ovarian follicles, whereas ovarian apoptosis is enhanced by IGF binding protein (52). Available research findings suggest that melatonin influences the process of steroidogenesis at the molecular level especially mRNA expression of CYP 11A and CYP17 gene which are the rate-limiting steps in this biosynthesis pathway (38).

### 5.4. Effect of melatonin on ovulation

Ovulation is a complex process by which a preovulatory follicle ruptures and releases a fertilizable oocyte into the oviductal lumen. This process occurs as a result of a dynamic interaction between the LH surge and local factors including

steroids, nitric oxide (NO), prostaglandins, and peptides in a time-dependent manner. The LH surge activates basic structural and biochemical variations leading to rupture of the graafian follicles, and, therefore, resulting in the exclusion of the oocyte with subsequent development of a corpus luteum (CL). Combined treatment of melatonin and progesterone (P) into women resulted in a reduction of LH secretion and finally block ovulation. It is also reported that such combined therapy led to an increase in circulating P without effecting FSH or inhibiting LH secretion (53). This effect might have mediated by the influence of melatonin on hypothalamic gonadotropin release via activation of the HPG axis (54). Nonetheless, the experiment showed that large follicles showed higher melatonin concentration as compared to smaller follicles therefore, strongly suggesting a positive correlation between follicular P and melatonin concentrations (31). A Higher concentration of circulatory melatonin during the preovulatory phase is suggested to be involved in the production of progesterone, consequently leading in luteinization, followed by successful ovulation (31). In this connection, research findings confirm that melatonin treatment (20 mg/kg body weight) resulted in a significant increment of prostaglandin E2 concentrations in gastric mucosa as well as esophageal tissue in rats (55-56). On the other hand, the inhibitory effect of melatonin on prostaglandin E2 in medial basal hypothalamus has also been documented by researchers (57). Here it becomes important to address that although the relationship between Prostaglandin E2 and melatonin received enough experimental pieces of evidence by the researcher even then the mechanism towards their exact involvement is still not properly established. Nevertheless, the process of ovulation appears very similar to a local inflammatory response because of the generation of both RNS and ROS (58). Mammalian ovulation is comparable to the inflammatory reaction as successful ovulation requires elevated follicular prostaglandin E2 levels which is one of the inflammatory components. Elevation in follicular prostaglandin levels activates proteolytic enzymes and enhancement of vascular permeability (58). Both endothelial NO synthase (eNOS) and

inducible NO synthase (iNOS) are present in the oocytes and thecal cells of the mouse (59). The major source of reactive oxygen species (ROS) appears to be inflammatory to cells including macrophages and neutrophils, as they are present in the ovary during ovulation. They generate tremendous numbers of free radicals leading to the induction of apoptosis in ovarian cells (6063). Melatonin, as well as its metabolites, is broad-spectrum antioxidants and free radical scavengers that quenches ROS as well as reactive nitrogen species (RNS) (64- 66). Elevated circulatory level of plasma melatonin of preovulatory follicles is presumed to protect GCs and the oocyte from free radicals stress and thus helping in the induction of ovulation (67-68).

### 5.5. Effect of melatonin in maintenance of Oocyte quality

Poor oocyte quality remains a profound problem for female infertility and hence contributes and participates in the process of fertilization. Oxidative stress, due to ROS generation during the ovulatory process within the follicle, may be a cause of poor oocytes quality (69). They accelerate apoptosis cause deterioration of cell membrane lipids and destroy DNA *etc.* (70). The balance between ROS production and the scavenging ability of antioxidants is an important factor during the process of oocyte maturation and till fertilization. Further, drugs that protect the oocyte and its surrounding feeder cells from damage, are of great importance and in this connection, melatonin is considered on the top among the basic pinealologist. In this review it is previously discussed that follicular fluid (FF) showed a significantly high level of endogenous melatonin] and its receptors in GCs suggesting a highly beneficial role of this indoleamine and neurohormone in the follicle at the molecular level (26-27,31,35). Melatonin administration (5 mg/kg body weight) enhances SOD activity and at physiological serum concentrations (1 nM), induces gene expression of all the three antioxidant enzymes (i.e., Cu, Zn-SOD, Mn-SOD, and GPx) (71- 72). Hence, melatonin positively induces both antioxidant enzyme activity as well as the expression of their corresponding genes. Therefore, it may be suggested that melatonin could become the medicine of choice for improving oocyte

quality for women who are unable to become pregnant because of poor oocyte quality. Melatonin also regulates the oocyte maturation capacity (73).

Accelerated action of maturation-inducing hormone for the formation of maturation- promoting factor was reported, following administration of melatonin (50–500 pg/mL) and germinal vesicle breakdown of oocytes. It exerts DNA methyltransferase inhibitory effects by masking target sequences and/or by blocking the active site of the enzyme (74). Melatonin's epigenetic efficacy has been proven in several diseases including cancer and hypertension (75-76). However, this miraculous neurohormone may also induce epigenetic changes in oocytes (77). These epigenetic modifications may result from the interaction of melatonin with nuclear melatonin receptors (78). The nuclear receptors of melatonin seem to show a functional role in DNA bending because it may result in the changes of DNA superstructure following epigenetic modifications by affecting nuclear melatonin receptors (79). Therefore, melatonin seems to be a mediator, to oocytes and interaction between environmental factors and the epigenetic inheritance system, for the environmental stimuli. Melatonin being a lipophilic molecule as well as an effective free radical scavenger in the culture medium supports not only mouse fertilization but also to the early development of embryonic tissue (80). Earlier it has been reported that melatonin supplementation (10 nM) consequences a positive effect on porcine embryo cleavage rates and blastocyst total cell numbers (81). Besides, melatonin in the culture medium improved the rate of development of thawed blastocysts with a higher hatching rate after 24 hours of culture (82). No negative effects of melatonin were observed on embryo development, at any of the concentrations 1 pM–100 tested, or even when administered in high doses during pregnancy (83-84).

### 6. MODULATION OF LUTEAL FUNCTION BY MELATONIN

The formation of the CL is initiated by a series of morphological and biochemical changes in cells of the theca interna and granulosa of the preovulatory follicle (85). Melatonin may act at the level of the ovary to modulate its luteal function. Not

only structural changes but genomic alterations lead to the terminal differentiation of follicular cells into P-producing luteal cells. Research reports deliver information regarding the increase in melatonin levels in the luteal phase compared with the follicular phase of the menstrual cycle (86-87). In humans, melatonin binding sites have been detected in GCs–luteal cells which consequently stimulate the secretion of P by human GCs or luteal cells (26, 31, 35). This, in turn, remarkably increases mRNA expression of the LH (but not FSH) receptor in human GCs/luteal cells, while inhibited expression of GnRH therefore, the GnRH receptor indicating a direct involvement in the regulation of the luteal function (26,31,35). Further, this hormone also enhances hCG-stimulated P secretion from these cells, possibly by the elevated expression of the LH receptor. On the contrary, some reports showed no effects or negative effects of melatonin on P production in the growing and luteinized GCs (28, 29, 50, 89). Nonetheless, literature also suggests the inhibitory action of melatonin (1 ng/mL–100 mg/mL), P and cAMP secretion by GCs isolated from porcine ovaries (89). Interestingly, the effects of melatonin such as induction of mRNA expression for the LH receptor and supportive effect on GCs are more dominant than its inhibitory actions. Melatonin might rescue GCs from induced free radical cytotoxicity in long-term cultured cells by its direct and indirect antioxidative ability.

The essential effect of melatonin on endometrial morphology and embryo implantation has been well described by Dair and their associated research team (90). The reports demonstrated that the implantation rates and serum P levels were decreased in the pinealectomized rats, whereas the reduced serum P levels were restored to normal by daily melatonin injections (2 mg/kg). Nevertheless, melatonin also increases PRL secretion inhibits the oxytocin release from the rat's hypothalamic-neurohypophysial system, suggesting that the indoleamine is essential for maintaining P production and luteal function by making the best use of various mechanisms (91-92). Elevated melatonin in the luteal phase and early pregnancy may induce P production by luteal cells, which is necessary for a successful pregnancy.

## 7. PROTECTIVE EFFECT OF MELATONIN IN EMBRYO CULTURE

Oocytes and embryos face the stress of high concentrations of reactive oxygen species (ROS) during gamete and embryo culture in the Assisted reproductive technology (ART) technique. Nowadays, infertility research working on assisted reproductive technology (ART) has developed an intense interest in the effects of oxidative stress on the success rates of ART. The retrieval of the micro-environment for cultured gametes and embryos is taken as an essential determinant factor for fertilization and consequently successful implantation. The treatment of melatonin has been reported beneficial in previous *in vivo* studies when it was supplemented in corresponding culture media of bovine, murine, and porcine embryo development (93). Ranges of melatonin concentration between 10–5M and 10–11 M showed an increase in 8-celled embryo yield and hence an increased number of blastocysts and blastocyst hatching in embryos with a higher cleavage rate in bovine (94). Findings from the previous experiment lead to the hypothesis that lower concentration of melatonin has a significant impact on the in-vitro supplementation of embryo culture media with its higher impact on the development and quality of embryos and also reported to be less harmful than higher doses (94).

## 8. EFFECT OF MELATONIN IN SUCCESS RATE OF IVF

Improvement in the success rates of IVF remains a focus of infertility research. As discussed earlier regarding the increment in the reactive oxygen species during the process of assisted reproductive techniques like *in vitro* fertilization (IVF) which undesirably may cause cell death and hence negatively affects the success rate of IVF outcome (95). The participation of the important role of melatonin during the luteal phase is also reported in patients undergoing IVF treatment. In 25 women with luteal phase defect and between them 14 women given with 3 mg/day of melatonin calculated with time-dependent delivery to match the HCG trigger time throughout the luteal phase it was found in a prospective study that melatonin supplementation significantly increased progesterone levels (11.0



ng/mL vs 8.9 ng/mL,  $p < 0.05$ ) (96). A survey-based on for IVF-embryo transfer suggests that melatonin concentration is higher in the fluid of large follicles than in the smaller follicles (48) Also, the fertilization success of the women with a low fertilization rate (%50%) in the prior IVF-embryo transfer cycle was improved by melatonin treatment compared to the prior IVF-embryo melatonin, because of being a strong antioxidant molecule, may enhance the improvement of egg quality which further can ensure the increased success rate for IVF and could be a good *in vitro* drug for the ARTs infertility clinics (97).

## 9. CONCLUSION

Melatonin has received a lot of attention in recent years with a major focus on its free radical scavenging capacity. These findings have widened the understanding of its pleiotropic physiological effects. Among these physiological actions, few are receptor-dependent whereas others are receptor-independent. The presence of melatonin has been reported in FF which might be taken up from the circulation after being released from the pineal gland. During the entire process of ovulation free radicals are produced. Melatonin because of its, antioxidative and free radical scavenging property scavenge produced free radicals; diminishes oxidative stress thereby participating in ovulation, oocyte maturation, and development of an embryo. Findings also indicate its potential role in ART and hence participating as an antioxidant for women having IVF with the aim of possible improvement of egg quality. The safety evaluation of *in vitro* and *in vivo* melatonin treatment has been broadly demonstrated by various researchers, suggesting without any unfavorable impact. However, before adopting the administration of melatonin it becomes important to further evaluate pharmacological consequences and therapeutic potentials. On a closing note, it is expected that melatonin or its novel analog will certainly find a good place among reproductive medicine in the near future.

## 10. ACKNOWLEDGMENTS

The authors extend thank to Dr. Shail K Chabe for his intellectual encouragement and Siamak

TabiZadeh for his comprehensive editorial work and efforts.

## 11. REFERENCES

1. Carrillo-Vico A, Calvo JR, Abreu P, Lardone PJ, García-Mauriño S, Reiter RJ, Guerrero JM: Evidence of melatonin synthesis by human lymphocytes and its physiological significance: possible role as intracrine, autocrine, and/or paracrine. *FASEB J.*, 18(3), 537-9 (2004) DOI: 10.1096/fj.03-0694fje
2. Rai S, Haldar C: Melatonin ameliorates oxidative stress and induces cellular proliferation of lymphoid tissues of a tropical rodent, *Funambulus pennanti*, during reproductively active phase. *Protoplasma*, 250 (1), 21-32 (2011) DOI: 10.1007/s00709-011-0367-1
3. Bubenik GA: Gastrointestinal melatonin: localization, function, and clinical relevance. *Dig Dis Sci*, 47, 2336-2348 (2002) DOI: 10.1023/A:1020107915919
4. Rai S, Basheer M, Ghosh H, Acharya D, Hajam Y A: Melatonin attenuates free radical load and reverses histologic architect and hormone profile alteration in female rat: An *in vivo* study of pathogenesis of letrozole induced polycystic ovary. *Clin & Cellular Immunol*. 6,384 (2015) DOI: 10.4172/2155-9899.1000384
5. Sharkey KM, Eastman CI: Melatonin phase shifts circadian rhythms in a placebo-controlled stimulated night-work study. *Am J Physiol Regul Integr Comp Physiol*, 282(2), R454-R463 (2002) DOI: 10.1152/ajpregu.00135.2001
6. Luboshitzky R, Yanaid D, Shen-Orr Z, Israilei E, Herer P, Lavie P: Daily and

- seasonal variations in the concentration of melatonin in the human pineal gland. *Brain Res, Bull.* 47 (3), 271-276 (1998)  
DOI: 10.1016/s0361-9230(98)00105-1
7. Duffy JF, Dijk DJ, Hall EF, Czeisle CA: Relationship of endogenous circadian melatonin and temperature rhythms to self-reported preference for morning or evening activity in young and old people. *J Invest Med* , 47, 141-150 (1999)  
<http://epubs.surrey.ac.uk/id/eprint/822581>
8. Haldar C, Hausseler D, Gupta D: Effect of pineal gland on circadian rhythmicity of colony forming units for granulocytes and macrophages (CFU-GM) from rat bone marrow cell –culture. *J Pineal Res*, 12, 79-83 (1992)  
DOI: 10.1111/j.1600-079X.1992.tb00030.x
9. Ghosh H, Rai S, Basheer M, Hajam YA: Receptor Cross Talk and Interplay between Melatonin and Ovarian Thyroid Axis in a Letrozole Induced Polycystic (Pco) Rat. *J Clin Cell Immunol*, 9, 552. (2018)  
DOI: 10.4172/2155-9899.1000552
10. Armstrong S M: Melatonin : the internal Zeirgeber of mammals. In *Pineal Research Review*, 7, 158-202 Ed. R. J. Reiter Alan R. Liss, New York (1989)
11. Rai S, Haldar C: Pineal Control of immune status and hematological changes in blood and bone marrow of male squirrels (*Funambulus pennanti*) during their reproductively inactive phase. *Comp Biochem Physiol C Toxicol Pharmacol* ,136, 319-328 (2003)  
DOI: 10.1016/j.cca.2003.10.008
12. Reiter RJ, Calvo JR, Karbownik M, Qi W, Tan D X: Melatonin and its relation to immune system and inflammation. *Ann N Y Acad Sci* , 917, 376-386 (2000)  
DOI: 10.1111/j.1749-6632.2000.tb05402.x
13. Tan DX, Reiter RJ , Manchester LC, Yan MT, El-Sawi M, Sainz RM, Mayo JC, Kohen R, Allegra M, Hardeland R: Chemical and Physical Properties and Potential Mechanisms: Melatonin as a Broad Spectrum Antioxidant and Free Radical Scavenger. *Current Topics in Medicinal Chemistry*, 2, 181-197 (2002)  
DOI: 10.2174/1568026023394443
14. Hajam YA, Rai S: Melatonin and insulin modulates the cellular biochemistry, histoarchitecture and receptor expression during hepatic injury in diabetic rats. *Life Sci*, 239,117046. (2019)  
DOI: 10.1016/j.lfs.2019.117046
15. YA Hajam, S Rai, A Roy, M Basheer, H Ghosh: Repossession of brain complications in a streptozotocin induced diabetic rat by exogenous melatonin administration. *Int. J. Zool. Res*, 13, 64-73 (2017)  
DOI: 10.3923/ijzr.2017.64.73
16. S Rai, YA Hajam, M Basheer, H Ghosh: Biochemical and histopathological inflections in hepato-renal tissues of streptozotocin (STZ) induced diabetic male rats: Impact of exogenous melatonin administration. *J. Clin. Res. Bioeth*, 7, 10.4172 (2016)  
DOI: 10.4172/2155-9627.1000290
17. Cecon E, Oishi A, Jockers R: Melatonin receptors: molecular pharmacology and signaling in the context of system bias. *Brit J Pharmacol*, 175, 3263–80 (2018)  
DOI: 10.1111/bph.13950

18. Slominski A, Fischer TW, Zmijewski MA, Wortsman J, Semak I, Zbytek B, Slominski RM, Tobin DJ: On the role of melatonin in skin physiology and pathology. *Endocrine*, 27(2), 137-14 (2005)  
DOI: 10.1385/ENDO:27:2:137
19. Carrillo-Vico A, Calvo JR, Abreu P, Lardone PJ, GarcíaMauriño S, Reiter RJ, Guerrero JM: Evidence of melatonin synthesis by human lymphocytes and its physiological significance: possible role as intracrine, autocrine, and/or paracrine substance. *FASEB J*, 18, 537-553 (2004)  
DOI: 10.1096/fj.03-0694fje
20. Cardinali DP, Rosner JM. Metabolism of serotonin by the rat retina *in vitro*. *J Neurochem*, 18 (9), 1769-1770 (1971)  
DOI: 10.1111/j.1471-4159.1971.tb03752.x
21. Kim MK, Park EA, Kim HJ, Choi WY, Cho JH, Lee WS, Cha KY, Kim YS, Lee DR, Yoon TK: Does supplementation of in-vitro culture medium with melatonin improve IVF outcome in PCOS? *Reprod Biomed*, 26 (1), 22-9 (2013)  
DOI: 10.1016/j.rbmo.2012.10.007
22. Waldhauser F, Ehrhart B, Förster E: Clinical aspects of the melatonin action: impact of development, aging, and puberty, involvement of melatonin in psychiatric disease and importance of neuroimmunoendocrine interactions. *Experientia*, 49, 671–681 (1993)  
DOI: 10.1007/BF01923949
23. García MJ, Hoyos AM, Carballo AM, García JM, López EN, Fernández JU: Puberty and melatonin. *An Esp Pediatr*, 57(2), 121-126 (2002)
24. Aleandri V, Spina V, Ciardo A: The role of the pineal body in the endocrine control of puberty. *Minerva Ginecol*, 49(1-2):43-48 (1997)
25. Silman R: Melatonin and the human gonadotrophin-releasing hormone pulse generator. *J Endocrinol*, 128(1), 7- 11 (1991)  
DOI: 10.1677/joe.0.1280007
26. Cohen HN, Hay ID, Annesley TM, Beastall GH, Wallace AM, Spooner R, Thomson JA, Eastwold P, Klee GG: Serum immunoreactive melatonin in boys with delayed puberty. *Clin Endocrinol (Oxf)*, 17 (5): 517-521(1982)  
DOI: 10.1111/j.1365-2265.1982.tb01623.x
27. Rönnerberg L, Kauppila A, Leppäluoto J, Martikainen H, Vakkuri O: Circadian and seasonal variation in human preovulatory follicular fluid melatonin concentration. *J Clin Endocrinol Metab*, 71(2), 493-496 (1990)  
DOI: 10.1210/jcem-71-2-493
28. Brzezinski A, Seibel MM, Lynch HJ, Deng MH, Wurtman RJ: Melatonin in human preovulatory follicular fluid. *J Clin Endocrinol Metab*, 64(4), 865-867 (1987)  
DOI: 10.1210/jcem-64-4-865
29. Nakamura Y, Tamura H, Takayama H, Kato H: Increased endogenous level of melatonin in preovulatory human follicles does not directly influence progesterone production. *Fert and Ster*, 80(4), 1012-1016 (2003)  
DOI: 10.1016/S0015-0282(03)01008-2
30. Murayama T, Kawashima M, Takahashi T, Yasuoka T, Kuwayama T, Tanaka K: Direct action of melatonin on hen ovarian granulosa cells to lower responsiveness

- to luteinizing hormone. *Proc Soc Exp Biol Med*, 215(4),386-392 (1997)  
DOI: 10.3181/00379727-215-44148
31. Woo MM, Tai CJ, Kang SK, Nathwani PS, Pang SF, Leung PC: Direct action of melatonin in human granulosa-luteal cells. *J. Clin. Endocrinol. Metab*, 86(10):4789-4797 (2001)  
DOI: 10.1210/jcem.86.10.7912
32. Tamura H, Nakamura Y, Takiguchi S, Kashida S, Yamagata Y, Sugino N, Kato H: Melatonin directly suppresses steroid production by preovulatory follicles in the cyclic hamster. *J Pineal Res*, 25(3), 135-141 (1998)  
DOI: 10.1111/j.1600-079X.1998.tb00551.x
33. Webley GE, Leidenberger F: The circadian pattern of melatonin and its positive relationship with progesterone in women. *J Clin Endocrinol Metab*, 63(2),323-328. (1986)  
DOI: 10.1210/jcem-63-2-323
34. Brzezinski A, Fibich T, Cohen M, Schenker JG, Laufer N: Effects of melatonin on progesterone production by human granulosa lutein cells in culture. *Fert Ster*, 58(3), 526-529 (1992)  
DOI: 10.1016/S0015-0282(16)55257-1
35. Yie SM, Brown GM, Liu GY, Collins JA, Daya S, Hughes EG, Foster WG, Younglai EV: Melatonin and steroids in human pre-ovulatory follicular fluid: seasonal variations and granulosa cell steroid production. *Hum reprod*, 10(1),50-55 (1995)  
DOI: 10.1093/humrep/10.1.50
36. Yie SM, Niles LP, Younglai EV: Melatonin receptors on human granulosa cell membranes. *J Clin Endocrinol Metab*, 80(5),1747-1749 (1995)  
DOI: 10.1210/jcem.80.5.7745030
37. Niles LP, Wang J, Shen L, Lobb DK, Younglai EV: Melatonin receptor mRNA expression in human granulosa cells. *Mol Cell Endocrinol*, 156(1-2),107-110 (1999)  
DOI: 10.1016/s0303-7207(99)00135-5
38. Soares JM, Masana MI, Erşahin Ç, Dubocovich ML: Functional melatonin receptors in rat ovaries at various stages of the estrous cycle. *J Pharmacol Exp Ther*, 306(2),694-702 (2003)  
DOI: 10.1124/jpet.103.049916
39. Soares JM, Simoes MJ, Oshima CT, Mora OA, De Lima GR, Baracat EC: Pinealectomy changes rat ovarian interstitial cell morphology and decreases progesterone receptor expression. *Gynecol Endocrinol*, 17(2),115-123 (2003)  
DOI: 10.1080/gye.17.2.115.123
40. Mor M, Plazzi PV, Spadoni G, Tarzia G: *Melatonin. Curr Med Chem*, 6,501–18 (1999)
41. Rivara S, Lorenzi S, Mor M, Plazzi PV, Spadoni G, Bedini A, Tarzia G: Analysis of structure– activity relationships for MT2 selective antagonists by melatonin MT1 and MT2 receptor models. *J. Med. Chem*, 48(12),4049-4060 (2005)  
DOI: 10.1021/jm048956y
42. von Gall C, Stehle JH, Weaver DR: Mammalian melatonin receptors: molecular biology and signal transduction. *Cell Tissue Res*, 309(1):151-162 (2002)  
DOI: 10.1007/s00441-002-0581-4



43. Reppert SM: Melatonin receptors: molecular biology of a new family of G protein-coupled receptors. *J Biol Rhythms*, 12(6):528-31 (1997)  
DOI: 10.1177/074873049701200606
44. Vanecek J: Cellular mechanisms of melatonin action. *Physiol Rev*, 78(3):687-721(1998)  
DOI: 10.1152/physrev.1998.78.3.687
45. Itoh MT, Ishizuka B, Kuribayashi Y, Amemiya A, Sumi Y: Melatonin, its precursors, and synthesizing enzyme activities in the human ovary. *Molecular Human Reproduction*, 5(5), 402-408 (1999).  
DOI: 10.1093/molehr/5.5.402
46. Wurtman RJ, Axelrod J, Potter LT: The uptake of H3-melatonin in endocrine and nervous tissues and the effects of constant light exposure. *Journal of Pharmacology and Experimental Therapeutics*, 143,314-318 (1964)
47. Webb R, Nicholas B, Gong JG, Campbell BK, Gutierrez CG, Garverick HA, Armstrong DG: Mechanisms regulating follicular development and selection of the dominant follicle. *Reproduction (Cambridge, England) Supplement*, 61:71-90 (2003)  
DOI: 10.1530/biosciprocs.5.007 .
48. Wu CS, Leu SF, Yang HY, Huang BM: Melatonin inhibits the expression of steroidogenic acute regulatory protein and steroidogenesis in MA-10 cells. *J Andrology*, 22(2),245–54 (2001)  
DOI: 10.1002/j.1939-4640.2001.tb02177.x
49. Schaeffer HJ, Sirotkin AV: Melatonin and serotonin regulate the release of insulin-like growth factor-I, oxytocin and progesterone by cultured human granulosa cells. *Exp Clin Endocrinol Diabetes*, 105(02),109-112(1997)  
DOI: 10.1055/s-0029-1211736
50. Picinato MC, Hirata AE, Cipolla-Neto J, Curi R, Carvalho CR, Anhe GF, Carpinelli AR: Activation of insulin and IGF-1 signaling pathways by melatonin through MT1 receptor in isolated rat pancreatic islets. *J Pineal Res*, 44(1), 88-94 (2008)  
DOI: 10.1111/j.1600-079X.2007.00493.x
51. Poretsky L: The insulin-related ovarian regulatory system in health and disease. *Endocr Rev*, 20 (4):535-582 (1999)  
DOI: 10.1210/edrv.20.4.0374
52. Chun SY, Billig H, Tilly JL, Furuta I, Tsafiri A, Hsueh AJ: Gonadotropin suppression of apoptosis in cultured preovulatory follicles: mediatory role of endogenous insulin-like growth factor I. *Endocrinology*, 135(5), 1845-1853 (1994)  
DOI: 10.1210/en.135.5.1845
53. Voordouw BC, Euser R, Verdonk RE, Alberda BT, de Jong FH, Drogendijk AC, Fauser BC, Cohen M: Melatonin and melatonin-progestin combinations alter pituitary-ovarian function in women and can inhibit ovulation. *J Clin Endocrinol Metab*, 74(1), 108-117 (1992)  
DOI: 10.1210/jcem.74.1.1727807
54. Roy SK, Greenwald GS: *In vitro* steroidogenesis by primary to antral follicles in the hamster during the periovulatory period: effects of follicle-stimulating hormone, luteinizing hormone, and prolactin. *Biol Reprod*, 37(1), 39-46 (1987)  
DOI: 10.1095/biolreprod37.1.39
55. Cabeza J, Alarcón-de-la-Lastra C, Jiménez D, Martín MJ, Motilva V:

- Melatonin modulates the effects of gastric injury in rats: role of prostaglandins and nitric oxide. *Neurosignals*, 12(2), 71-7 (2003)  
DOI: 10.1159/000071816
56. Konturek SJ, Zayachkivska O, Havryluk XO, Brzozowski T, Sliwowski Z, Pawlik M, Konturek PC, Czesnikiewicz-Guzik M, Gzhegotsky MR, Pawlik WW: Protective influence of melatonin against acute esophageal lesions involves prostaglandins, nitric oxide and sensory nerves. *J Physiol Pharmacol*, 58(2), 361-377 (2007)
  57. Bettahi I, Guerrero JM, Reiter RJ, Osuna C. Physiological concentrations of melatonin inhibit the norepinephrine-induced activation of prostaglandin E2 and cyclic AMP production in rat hypothalamus: A mechanism involving inhibition of nitric oxide synthase. *J Pineal Res*, 25(1), 34-40 (1998)  
DOI: 10.1111/j.1600-079X.1998.tb00383.x
  58. Espey LL: Current status of the hypothesis that mammalian ovulation is comparable to an inflammatory reaction. *Biol Reprod*, 50(2), 233-238 (1994)  
DOI: 10.1095/biolreprod50.2.233
  59. Mitchell LM, Kennedy CR, Hartshorne GM: Expression of nitric oxide synthase and effect of substrate manipulation of the nitric oxide pathway in mouse ovarian follicles. *Hum Reprod*, 19(1), 30 -40 (2004)  
DOI: 10.1093/humrep/deh032
  60. Nakamura Y, Smith M, Krishna A, Terranova PF: Increased number of mast cells in the dominant follicle of the cow: relationships among luteal, stromal, and hilar regions. *Biol Reprod*, 37(3),546-549 (1987)  
DOI: 10.1095/biolreprod37.3.546
  61. Brännström M, Mayrhofer G, Robertson SA: Localization of leukocyte subsets in the rat ovary during the periovulatory period. *Biology of Reproduction*, 48(2), 277-286 (1993)  
DOI: 10.1095/biolreprod48.2.277
  62. Gupta RK, Miller KP, Babus JK, Flaws JA: Methoxychlor inhibits growth and induces atresia of antral follicles through an oxidative stress pathway. *Toxicol Sci*, 93(2),382-389 (2006)  
DOI: 10.1093/toxsci/kfl052
  63. Korzekwa AJ, Okuda K, Woclawek-Potocka I, Murakami S, Skarzynski DJ: Nitric oxide induces apoptosis in bovine luteal cells. *J Reprod Dev*, 52(3), 353-361 (2006)  
DOI: 10.1262/jrd.17092
  64. Tripathi A, Prem Kumar Karuppana V, Pandey Ashutosh N, Khatun Shabana, Kirti Mishra Surabjhi, Shrivatav Tulsidas G, Chaube Shail K : Melatonin protects against clomiphene citrate-induced generation of hydrogen peroxide and morphological apoptotic changes in rat eggs. *Eur J of Pharmacology*, 667 (1-3), 419-424 (2011)  
DOI: 10.1016/j.ejphar.2011.06.005
  65. Reiter RJ, Tan DX, Maldonado MD: Melatonin as an antioxidant: physiology versus pharmacology. *J Pineal Res*, 39(2), 215-216 (2005)  
DOI: 10.1111/j.1600-079X.2005.00261.x
  66. Tan DX, Reiter RJ, Manchester LC, Yan MT, El-Sawi M, Sainz RM, Mayo JC, Kohen R, Allegra MC, Hardeland R: Chemical and physical properties and

- potential mechanisms: melatonin as a broad spectrum antioxidant and free radical scavenger. *Current Topics in Medicinal Chemistry*, 2(2),181-197 (2002)  
DOI: 10.2174/1568026023394443
67. Reiter RJ , Tan DN, Gitto E, Sainz RM, Mayo JC, Leon J, Manchester LC, Vijayalaxmi, Kilic E, Kilic U: Pharmacological utility of melatonin in reducing oxidative cellular and molecular damage. *Polish J Pharmacol* 56(2),159-170 (2004)
  68. Tan DX, Manchester LC, Terron MP, Flores LJ, Reiter RJ: One molecule, many derivatives: A never-ending interaction of melatonin with reactive oxygen and nitrogen species? *J Pineal Res*, 42(1),28-42 (2007)  
DOI: 10.1111/j.1600-079X.2006.00407.x
  69. Agarwal A, Gupta S, Sharma RK: Role of oxidative stress in female reproduction. *Reprod Biol Endocrinol*, 3(1),28 (2005)  
DOI: 10.1186/1477-7827-3-28
  70. Kowaltowski AJ, Vercesi AE: Mitochondrial damage induced by conditions of oxidative stress. *Free Radic Biol Med*, 26(3-4), 463-471 (1999)  
DOI: 10.1016/s0891-5849(98)00216-0
  71. Liu F, Ng TB: Effect of pineal indoles on activities of the antioxidant defense enzymes superoxide dismutase, catalase, and glutathione reductase, and levels of reduced and oxidized glutathione in rat tissues. *Biochem Cell Biol*, 78(4),447-453 (2000)  
DOI: 10.1139/o00-018
  72. Mayo JC, Sainz RM, Antoli I, Herrera F, Martin V, Rodriguez C: Melatonin regulation of antioxidant enzyme gene expression. *Cell Mol Life Sci*,59(10),1706-1713 (2002)  
DOI: 10.1007/pl00012498
  73. Adriaens I, Jacquet P, Cortvrindt R, Janssen K, Smitz J: Melatonin has dose-dependent effects on folliculogenesis, oocyte maturation capacity and steroidogenesis. *Toxicology*, 228(2-3),333-343 (2006)  
DOI: 10.1016/j.tox.2006.09.018
  74. Chatteraj A, Bhattacharyya S, Basu D, Bhattacharya S, Bhattacharya S, Maitra SK: Melatonin accelerates maturation inducing hormone (MIH): induced oocyte maturation in carps. *Gen Comp Endocrinol*, 140(3),145-155 (2005)  
DOI: 10.1016/j.ygcen.2004.10.013
  75. Ya Li, Sha Li, Yue Zhou, Xiao Meng, Jiao-Jiao Zhang, Dong-Ping Xu, Hua-Bin Li: Melatonin for the prevention and treatment of cancer. *Oncotarget*, 13; 8(24): 39896–39921 (2017)  
DOI: 10.18632/oncotarget.16379
  76. Irmak MK, Sizlan A: Essential hypertension seems to result from melatonin-induced epigenetic modifications in area postrema. *Medical Hypotheses*, 66(5):1000-1007 (2006)  
DOI: 10.1016/j.mehy.2005.10.016
  77. Korkmaz A, Reiter RJ: Epigenetic regulation: a new research area for melatonin?. *J Pineal Res*, 44(1),41-44 (2008)  
DOI: 10.1111/j.1600-079X.2007.00509.x
  78. Missbach M, Jagher B, Sigg I, Nayeri S, Carlberg C, Wiesenberger I: Thiazolidine diones, specific ligands of the nuclear receptor retinoid Z receptor/retinoid acid receptor-related orphan receptor alpha

- with potent antiarthritic activity. *J Biol Chem*, 271(23),13515-13522 (1996)  
DOI: 10.1074/jbc.271.23.13515
79. Smirnov AN: Nuclear melatonin receptors. *Biochemistry (Mosc)*, 66(1),19-26 (2001)  
DOI: 10.1023/a:1002821427018
80. Ishizuka B, Kuribayashi Y, Murai K, Amemiya A, Itoh MT: The effect of melatonin on *in vitro* fertilization and embryo development in mice. *J Pineal Res*, 28(1),48-51 (2000)  
DOI: 10.1034/j.1600-079x.2000.280107.x
81. Rodriguez-Osorio N, Kim IJ, Wang H, Kaya A, Memili E: Melatonin increases cleavage rate of porcine preimplantation embryos *in vitro*. *J Pineal Res*, 43(3),283-8 (2007)  
DOI: 10.1111/j.1600-079X.2007.00475.x
82. Abecia JA, Forcada F: Zuniga O: The effect of melatonin on the secretion of progesterone in sheep and on the development of ovine embryos *in vitro*. *Veterinary Research Communications*. 26 (2),151-8 (2002)  
DOI: 10.1023/A:1014099719034
83. McElhinny AS, Davis FC, Warner CM: The effect of melatonin on cleavage rate of C57BL/6 and CB A/Ca preimplantation embryos cultured *in vitro*. *J Pineal Res*, 21(1),44-8 (1996)  
DOI: 10.1111/j.1600-079X.1996.tb00269.x
84. Jahnke G, Marr M, Myers C, Wilson R, Travlos G, Price C: Maternal and developmental toxicity evaluation of melatonin administered orally to pregnant Sprague-Dawley rats. *Toxicol Sci*, 50(2),271-9 (1999)
- DOI: 10.1093/toxsci/50.2.271
85. Richards JS, Russell DL, Ochsner S, Espey LL: Ovulation: new dimensions and new regulators of the inflammatory-like response. *Annu Rev Physiol*, 64(1), 69-92 (2002)  
DOI: 10.1146/annurev.physiol.-64.081501.131029
86. Wetterberg L, Arendt J, Paunier L, Sizonenko PC, Donselaar WV, Heyden T: Human serum melatonin changes during the menstrual cycle. *The Journal of Clinical Endocrinology & Metabolism*, 1976; 42(1), 185-188.  
DOI: 10.1210/jcem-42-1-185
87. Brun J, Claustrat B, David M: Urinary melatonin, LH, oestradiol, progesterone excretion during the menstrual cycle or in women taking oral contraceptives *Acta Endocrinol (Copenh)*, 116(1),145-149 (1987)  
DOI: 10.1530/acta.0.1160145
88. Bódis J, Koppan M, Kornya L, Tinneberg HR, Török A: Influence of melatonin on basal and gonadotropin-stimulated progesterone and estradiol secretion of cultured human granulosa cells and in the superfused granulosa cell system. *Gynecol Obstet Invest*, 52(3),198-202 (2001)  
DOI: 10.1159/000052973
89. Sirotkin AV: Direct influence of melatonin on steroid, Nona peptide hormones, and cyclic nucleotide secretion by granulosa cells isolated from porcine ovaries. *J Pineal Res*, 17(3), 112-117 (1994)  
DOI: 10.1111/j.1600-079X.1994.tb00121.x
90. Dair EL, Simoes RS, Simões MJ, Romeu LR, Oliveira-Filho RM, Haidar MA,



- Baracat EC, Soares JM. Effects of melatonin on the endometrial morphology and embryo implantation in rats. *Fertil Steril*, 89(5 Suppl),1299-1305 (2008)  
DOI: 10.1016/j.fertnstert.2007.03.050
91. Zisapel N: Melatonin–dopamine interactions: from basic neurochemistry to a clinical setting. *Cell Mol Neurobiol* , 21(6),605-616 (2001)  
DOI: 10.1023/A:1015187601628
92. Juszczak M, Stempniak B: Melatonin inhibits the substance P-induced secretion of vasopressin and oxytocin from the rat hypothalamo-neurohypophyseal system: *in vitro* studies. *Brain Res Bull*, 59(5), 393-397 (2003)  
DOI: 10.1016/s0361-9230(02)00942-5
93. Choi J, Park SM, Lee E, Kim JH, Jeong YI, Lee JY, Park SW, Kim HS, Hossein MS, Jeong YW, Kim S, Hyun SH, Hwang WS: Anti-apoptotic effect of melatonin on preimplantation development of porcine parthenogenetic embryos. *Mol Reprod Dev* 75(7):1127-1135 (2008)  
DOI: DOI: 10.1002/mrd.20861
94. Wang F, Tian X, Zhou Y, Tan D, Zhu S, Dai Y, Liu G: Melatonin improves the quality of *in vitro* produced (IVP) bovine embryos: implications for blastocyst development, cryotolerance, and modifications of relevant gene expression. *PIOS ONE*, 9(4):e93641. (2014)  
DOI: 10.1371/journal.pone.0093641
95. Paszkowski T, Clarke RN: Fertilization and early embryology: Antioxidative capacity of preimplantation embryo culture medium declines following the incubation of poor quality embryos. *Hum Reprod*. 11(11):2493-2495 (1996)  
DOI: 10.1093/oxfordjournals.-humrep.a019146
96. Tamura H, Takasaki A, Miwa I, Taniguchi K, Maekawa R, Asada H, Taketani T, Matsuoka A, Yamagata Y, Shimamura K, Morioka : Oxidative stress impairs oocyte quality and melatonin protects oocytes from free radical damage and improves fertilization rate. *J Pineal Res*, 44(3), 280-287 (2008)  
DOI: 10.1111/j.1600-079X.2007.00524.x
97. Fernando S Wallace ME Vollenhoven B Lolatgis N Hope N Wong M Lawrence M Lawrence A Russell C Leong K, P Rombauts L: Melatonin in Assisted Reproductive Technology: A Pilot Double-Blind Randomized Placebo-Controlled Clinical Trial. *Front Endocrinol(Lausanne)*, 9, 545 (2018)  
DOI: 10.3389/fendo.2018.00545

**Abbreviations:** SCN: Suprachiasmatic nucleus., LD: Light/dark cycle, LH: Luteinizing hormone, PCOS: Polycystic ovary syndrome , RZR: Retinoid Z receptor, ROR: Retinoic acid orphan receptor, MT1: Melatonin receptor 1, MT2: Melatonin receptor 2, cAMP: Cyclic adenosine 3050 monophosphate , FF: Follicular fluid , GC: Granulosa cell , FSH: Follicle stimulating hormone, IVF: In-vitro fertilization, NAT: N-acetyltransferase, HIOMT: Hydroxyindole-o-methoxytryptamine, E: Estrogen, P450scc: P450 side-chain cleavage enzyme , C17: P450 17a-hydroxylase,, P: Progesterone , E2: Estradiol , CYP: Cytochrome P, hCG: Human Chorionic gonadotrophin , IGF: Insulin like growth factor, PI3 K/AKT: Phosphatidylinositol 3- kinase/ protein kinase B , MEK: Mitogen activated protein kinase, ERK: Extracellular signaling regulated kinase, NO: Nitric oxide , ROS:

## Modulation of human ovarian functions by melatonin

Reactive oxygen species, GPx: Glutathione peroxidase , RNS: Reactive nitrogen species , CL: Corpus luteum , eNOS: Endothelial NO synthase , iNOS: Inducible NO synthase (iNOS), CL: Corpus leutum, PR: Progesterone receptor , GnRH: Gonadotropin regulatory hormone , PRL: Prolactin, PGF2a: ProstaglandinF-2a , LPO: Lipid peroxide , SOD: Superoxide dismutase , CAT: Catalase, ERa: Estrogen receptor a , ART: Assited Reproductive technology

**Key Words:** Pineal gland, Melatonin, Reproduction, Ovary, *In vitro* fertilization, IVF, Review

**Send correspondence to:** Seema Rai, Department of Zoology, Guru Ghasidas Vishwavidyalaya, Bilaspur Chhattisgarh, India, Tel: 07752-260472, Fax:,07752-260468 E-mail: seemarai.72@ggu.ac.in