

Bisphenol A: an endocrine-disruptor compound that modulates the immune response to infections

Victor Hugo Del Rio Araiza¹, Mariana Segovia-Mendoza², Karen Elizabeth Nava Castro³, Samira Muñoz Cruz⁴, Karina Chavez Rueda⁴, Carmen Tlazoelteotl Gomez de Leon⁵, Jorge Morales Montor⁶

¹*Departamento de Parasitología, Facultad de Medicina Veterinaria y Zootecnia, Universidad Nacional Autónoma de México, CP 04510, Ciudad de México, México,* ²*Departamento de Farmacología, Facultad de Medicina, Universidad Nacional Autónoma de México, CP 04510, Ciudad de México, México,* ³*Laboratorio de Genotoxicología y Mutagenesis Ambiental, Departamento de Ciencias Ambientales, Centro de Ciencias de la Atmosfera, Universidad Nacional Autónoma de México CP 04510, Ciudad de México, México,* ⁴*Unidad de Investigación Médica en Enfermedades Infecciosas y Parasitarias, Centro Médico Nacional Siglo XXI, Instituto Mexicano del Seguro Social, 06720, Cd de México, México.* ⁵*UIM en Inmunología, Hospital de Pediatría, CMN S XXI, IMSS, 06720, Cd de México, México.* ⁶*Departamento de Inmunología, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, 04510, Cd de México, México*

TABLE OF CONTENT

1. Abstract
2. Endocrine Disrupting Compounds (EDCS)
3. Bisphenol A (BPA)
4. BPA effects on the immune system
 - 4.1. BPA effects on innate immune cells
 - 4.1.1. Macrophages
 - 4.1.2. Dendritic cells
 - 4.1.3. Granulocytes
 - 4.1.4. Mast cells
5. BPA effects on adaptive immune cells
 - 5.1. T lymphocytes
 - 5.2. B lymphocytes and plasmatic cells
6. Effects of BPA on infections
 - 6.1. Virus
 - 6.2. Bacteria
 - 6.3. Parasites
7. Conclusions
8. Acknowledgments
9. References

1. ABSTRACT

Bisphenol A (BPA) is an endocrine-disruptor compound that exhibits estrogenic activity. BPA is used in the production of materials such as

polycarbonate plastics, epoxy resins and dental sealants. Whereas, the endocrine modulating activity of BPA and its effects on reproductive health have

BPA contributes to the patho-physiology of infectious diseases




Figure 1. Chemical structure of several EDCs. a) Synthetic origin; Dichloro-Diphenyl-Trichloroethane (DDT), Bisphenol A (BPA), Bisphenol S (BPS), Tetrabromo Bisphenol A (TetrabromoBPA), Ethinylestradiol, Diethylstilbestrol (DES). b) Natural origin; Genistein, Coumestrol, Daidzein.

been widely studied, its effects on the function of the immune system are poorly characterized. This might be attributable to the different BPA doses used in a diversity of animal models. Moreover, most studies of the effect of BPA on the immune response are limited to *in vitro* and *in vivo* studies that have focused primarily on the impact of BPA on the number and proportion of immune cell populations, without evaluating its effects on immune function in response to an antigenic challenge or infectious pathogens. In this review, we discuss the current literature on the effects of BPA on the function of immune system that potentially increases the susceptibility to infections by the virtue of acting as a pro-inflammatory molecule. Thus, it appears that BPA, while by such an impact might be useful in the control of certain disease states that are helped by an inflammatory response, it can worsen the prognosis of diseases that are adversely affected by inflammation.

2. ENDOCRINE DISRUPTING COMPOUNDS (EDCS)

Endocrine disrupting compounds (EDCs) are substances that exist in the environment as a result of agricultural and industrial activity. Some

examples are dichlorodiphenyltrichloroethane (DDT), Bisphenol A (BPA), Bisphenol S (BPS), phthalates, among others, but they also can be found in pharmaceutical products (ethinylestradiol, diethylstilbestrol (DES)) or naturally in different plants such as soybean (phytoestrogens, genistein, daidzein, coumestrol) (Figure 1). EDCs may exhibit estrogenic, anti-estrogenic or anti-androgenic activity. In addition, these compounds are highly lipophilic and can be stored for prolonged periods of time in the adipose tissues. During pregnancy, the fetus can be exposed to these compounds through the placenta and at birth by the lactogenic route (1, 2).

Different studies have shown that, perinatal and neonatal exposure to EDCs, generate irreversible alterations in the reproductive axis and in the central nervous system of the offspring of different species. Although, the exposure to EDCs during the perinatal and neonatal stages can lead to the development of different diseases in the adult life, EDCs may affect health at any age (3). This fact suggests that early life exposure to EDCs can increase the predisposition to a host of disease conditions.

BPA contributes to the patho-physiology of infectious diseases




Figure 2. Schematic representation of E2 and BPA interaction with nuclear ERs. E2 or BPA interacts with nuclear ERs. BPA; Bisphenol A, E2; Estradiol; ER α or β ; Estrogen receptor alpha or beta. The interaction between BPA and ER will activate genes such as ERK and NF κ B, which will generate the production of proinflammatory cytokines and nitric oxide. BPA; Bisphenol A, E₂; Estradiol, ER; Estrogen Receptor, ERK; extracellular signal-regulated kinases; NF κ B; nuclear factor kappa-light-chain-enhancer of activated B cells, NO; Nitric Oxide.

3. BISPHENOL A (BPA)

BPA is widely used as a monomer in the production of polycarbonate plastics, epoxy resins and dental sealants (4). This compound can be released easily from these materials due to incomplete polymerization or hydrolysis of the polymers that contain it, which can occur when they are exposed to high temperatures, acidic conditions or enzymatic process. The main BPA exposure source in animals and humans is through out food and beverages that have been in contact with materials manufactured with BPA, which is detached from its matrix and it is ingested orally (5). BPA is classified as an EDC with estrogenic character since it can bind to nuclear estrogen receptors (ER α and ER β) and trigger signaling pathways, even when its affinity is lower (<1,000) than the endogenous ligand, 17 β -estradiol (Figure 2) (6). In addition, BPA also binds to the estrogen like G-coupled protein receptor (GPER), the estrogen related receptor gamma (ERR γ), the arylhydrocarbon receptor (AhR) (7), the thyroid hormone receptor (ThR) (8), the peroxisome proliferator-activated receptor gamma (PPAR γ) (9),

the glucocorticoid receptor and the thyroid hormone receptor (TR) (8–15). Despite the Food and Drug Administration (FDA) and the European Food Safety Agency (EFSA) calculated that the Tolerable Daily Intake of BPA is 50 μ g / kg / day, it has been estimated that exposure to BPA per food package is 0.185 μ g / kg / day in adults and up to 2.42 μ g / kg / day in children from 1 to 2 months of age (8). Besides, it has been demonstrated that BPA exposure at the tolerable concentrations or below is related to negative effects in the health of both, humans and rodents (13, 16, 17).

4. BPA EFFECTS ON THE IMMUNE SYSTEM

Different BPA effects have been reported on the immune system cells; however, they vary depending on whether they were performed as *in vivo* or *in vitro*. *In vivo*, the effects reported may seem contradictories, but they are not, since the reported effects depend on the animal species used, the dose, the administration route, the sex of the animal, the age, and the animal's development stage in which BPA is administered. Furthermore, many reports do

BPA contributes to the patho-physiology of infectious diseases




Figure 3. Effects of BPA on the cells of the immune system. The BPA effects in different immune cells are variable depending on the type of model, *in vivo* or *in vitro*, the animal species used, the dose, the administration route and the stage of development in which it is administered. Mo; Macrophages, DC; Dendritic cells, Eos; Eosinophils, No; Neutrophils, PC; Plasma cells, BL; B lymphocytes, ThL; T helper lymphocytes, Treg; T regulatory lymphocytes.

not consider that the immune response must be studied by challenging the immune components, so there is little information about the BPA effects on the immune response during an infectious process. In the next paragraphs, we summarize the reported BPA effect on all the immune cells reported to date. To make it more comprehensive, we divide them in innate and adaptive immune cells (Figure 3).

4.1. BPA effects on innate immune cells

4.1.1. Macrophages

Macrophages are one of the main phagocytic cells that play an important role in the maintenance of organism homeostasis. Furthermore, they express the two ER isoforms (ER α and ER β) whereby BPA could exert its effects (18, 19). On line with that, different studies have evaluated the BPA stimulatory effects on macrophages. Hong *et al.* (2004) reported that BPA at a concentration of 43 nM potentiated nitric oxide production (NO) in a murine macrophage cell line (RAW264), after lipopolysaccharide (LPS) exposure; while interferon-gamma (IFN- γ) production was not altered (20). Other experiments developed by Yamashita *et al.* (2005) described that BPA at a concentration of 0.1

μ M stimulated pro-inflammatory cytokines production such as IL-1, IL-6 and IL-12, also BPA treatment increased the co-stimulatory molecule CD86 expression in murine peritoneal macrophages (21). In concordance with this study, mouse peritoneal macrophages under M1 type conditions that were exposed to low concentrations of BPA ($> 1 \mu$ M) promotes polarization toward M1 subtype by the upregulation of IRF5 expression, as well as TNF- α , IL-6 and MOP-1; while the same BPA exposure to macrophages under M2 type conditions inhibits the M2 subtype polarization by the downregulation of IL-10 and TGF- β (22). In addition, a BPA analog (BPA-glycidyl-methacrylate (BisGMA)) stimulated Tumor Necrosis Factor alpha (TNF- α) production with a concomitant increase of surface molecules (CD11, CD14, CD40, CD45, CD54 and CD80) expression on a macrophage cell line (RAW264.7). Like BPA, the BisGMA induced the production of IL-1 β , IL-6, and NO, as well as the inducible nitric oxide synthase (iNOS) expression and raised the generation of reactive oxygen species (ROS) intra and extracellular in a dose-dependent manner (23).

The increment of IL-6 and TNF- α secretion stimulated by BPA (0.1 μ M) was also observed in a

human macrophage cell line (THP1), on the contrary, secretion of regulatory cytokines such as IL-10 and TGF- β was decreased by BPA treatment. Moreover, the treatment of an ER antagonist (ICI 182,780) reversed this cytokine production pattern, indicating that BPA effects are carried out by its binding to ERs (24). Interestingly, BPA can induce alternative macrophages activation (M2), reducing IL-6, IL-10 and IL-1 β production in macrophages derived from human peripheral blood monocytes (PBMCs), stimulated or not with LPS or IL-4. This work also reported that BPA treatment increased the IL-10 and decreased the IL-6 production in macrophages classically activated (M1) (25). Finally, a work of Yang *et al.* (2015) reported that BPA increased the production of NO and ROS in a dose-dependent manner in carp (*Cyprinus carpio*) macrophages (26).

On the other hand, inhibitory BPA effects on macrophages function have also been reported. Segura *et al.* (1999) evaluated the adhesion capacity of the peritoneal macrophages of rats after BPA treatment, this compound (10 nM) inhibited the macrophages adherence but it did not have effect on their viability (27). In another study, Kim and Jeong (2003) evaluated the effect of BPA on the production of NO, TNF- α and iNOS expression in mice peritoneal macrophages. BPA (50 μ M) did not affect the NO and TNF- α production. On the contrary, when LPS was used as stimuli, BPA treatment inhibited their production. In addition, BPA decreased the iNOS expression in a dose-effect dependent manner (28). Supporting this fact, Byun *et al.* (2005) indicated that mice peritoneal macrophages, obtained from mice injected with BPA (500 mg / kg / day) for 5 consecutive days during 4 weeks, and cultured *ex vivo* with LPS, had a marked reduction in TNF- α and NO production. A similar effect was observed in a macrophage culture treated with BPA (10 and 100 Mm) (29). The BPA cell inhibitory effect was also observed in RAW 264 macrophage cell line, where BPA exposure (100 μ M) inhibited IFN- β promoter activation after LPS stimulation (30). Moreover, this estrogenic compound also suppressed the NO production and NF κ B activation in the same cell line with the same stimulus. Of note, these effects were blocked by the estrogen receptor antagonist ICI182,780 (31).

In line with that, BPA (200 μ M) exposure in RAW 264.7 macrophage cell line also shown to inhibit the production of NO and induce apoptosis cell death (32). In summary, BPA alters macrophage functions by decreasing cytokine secretion, activating of M2 phenotype and stimulate the expression of adhesion molecules.

Finally, in a recent study Liu *et al.* (2020), report that BPA (100 μ g / L) generates immunotoxicity in macrophages from common carp (*Cyprinus carpio*), by increasing the expression of long non-coding RNAs (33). This is reflected in the alteration of some signaling pathways related to the immune response, such as: NF- κ B, Toll-like receptor, B-cell receptor and the Jak-STAT signalling pathway (33).

4.1.2. Dendritic cells

Dendritic cells (DCs) are the par excellence antigen presenting cells; they play a fundamental role in the beginning of the immunological response and its polarization and regulation. It has been reported that these cells also express ER α and ER β (34). Limited studies have investigated the BPA effects on DCs. Guo Y and cols. (2010), reported that treatment of BPA in DCs derived from PBMCs increased the chemokine ligand 1 (CCL1) expression, the IL-5, IL-10 and IL-13 production as well as the expression of the transcription factor GATA3 in the presence of TNF- α (35), demonstrating that the exposure to BPA alters the functions of human DCs by inducing preferentially a Th2 response. Furthermore, BPA affected DCs differentiation by increasing the class II major histocompatibility complex (MHC II) and CD69 expression (36). On the other hand, Švajger *et al.* (2016), indicated that BPA at a concentration of 50 μ M decreased the DCs endocytic capacity as well as the CD1a expression (an important protein mediating antigenic presentation) (37). BPA at low concentrations (1 nM) also increased the DCs density, however, the expression of DCs activation markers (HLA-DR and CD86) was decreased in human DCs derived from PMBCs (38). BPA had no effect neither in cultivated bone marrow precursors or in the proliferation of the DCs, regardless of concentration (39). The works previously described suggest that exposure to BPA affects human DCs function in a concentration dependent manner.

4.1.3. Granulocytes

Granulocytes are the most abundant innate immune cells. They are divided into neutrophils, which constitutes between 90 and 95% of their totality, eosinophils from 3 to 5% and basophils less than 1%. In the literature there are few reports about the BPA effect on these cells. In the case of neutrophils, Watanabe *et al.*, (2003) evaluated the effect of BPA on the neutrophilic differentiation induced by dimethylsulfoxide (DMSO) and granulocyte colony stimulating factor (G-CSF) in a leukemia cell line (HL-60). They reported that low doses of BPA (10^{-10} and 10^{-8} M) increased the expression of the CD18 integrin protein, the neutrophilic differentiation as well as superoxide production. Interestingly, the treatment with an estrogen receptor inhibitor (tamoxifen) in these cells, did not suppress the BPA effect, suggesting that all the effects above are mediated by an ER-independent pathway (40). Another BPA-related compound, tetrabromobisphenol A (TBBPA), also increased the ROS neutrophils production in a dose-dependent manner (41).

BPA exposure has also been found to affect eosinophils, which are key players in allergy and asthma pathogenesis. *In vivo* experiments, using mouse models of allergic asthma, described that perinatal BPA exposure enhanced eosinophilic inflammation and airway responsiveness (42), suggesting that BPA exposure can affect the disease during critical developmental periods. Similar results were observed when adult mice were sensitized with ovalbumin combined with BPA treatment, where the result of the treatment significantly increased eosinophil recruitment in the alveoli and submucosa of the airways and enhanced eosinophil-produced cytokine and chemokine levels (43). These studies suggest that early or late life exposure to BPA could enhance the severity of immune-mediated diseases such as allergic diseases and asthma.

4.1.4. Mast cells

Mast cells are resident tissue cells that play a key role in inflammatory and allergic processes (44). These cells are characterized by their high content of cytoplasmic granules containing preformed mediators that include the vasoactive amine histamine, proteases, as well as some

cytokines (45). Mast cells may be activated by a variety of stimuli through the numerous receptors on their surface. Upon activation, mast cells can release preformed as well as a several distinct newly synthesized mediators (44,45). The most studied mechanism of mast cell activation is the IgE-mediated response, which can be divided into two stages; in the first stage, IgE produced by plasma cells binds to the Fc γ RI on the mast cell surface, leading to a sensitized state. Later, upon a second encounter with the antigen, it can bind to the surface-bound IgE, which after crosslink of IgE triggers mast cell degranulation and release of mediators. Though BPA exposure has been linked to allergen sensitization, there is limited information concerning BPA effects on mast cell-response. In a two-generational study, O'Brien reported that perinatal BPA exposure through maternal diet (ranging from 50 ng to 50 mg / Kg diet) caused an increase in mast cell-derived leukotrienes, prostaglandin D₂, TNF- α and IL-13, suggesting greater activation of these cells (46). Further studies will be needed to evaluate the role of the BPA exposure on mast cell response and its association to allergic diseases.

5. BPA EFFECTS ON ADAPTATIVE IMMUNE CELLS

5.1. T lymphocytes

T lymphocytes (TL) are cells of the adaptive immune system, they can be divided into cytotoxic T lymphocytes (CTLs) or T helper lymphocytes (ThL), which can differentiate into Th1, Th2, Th9, Th17, Th33., based on their secreting cytokines pattern. These cells express different hormone receptors (47), thus sex steroids can regulate the TLs differentiation and their response (48, 49). Some studies have reported that BPA exerts its effects through out the binding to them with a consequent modulation of TL response and polarization (Th1, Th2, and Th17).

In an *ex vivo* model, Youn *et al.*, (2002) described that BPA induced cell proliferation in Concanavalin A (Con-A)-stimulated splenocytes derived from mice orally exposed to BPA during 4 weeks, with no in ThL, CTL subpopulation percentages. In addition, the IFN- γ and the IL-4

expression was induced and reduced, respectively (50). Moreover, in chicken egg lysozyme (HEL) model, Yoshino *et al.*, (2003) reported that BPA significantly increased the IFN- γ and IL-4 secretion, suggesting that BPA promoted the Th1 response. This cytokine secretion pattern was also reported in prenatally BPA-exposed mice, after being immunized with HEL in the adulthood, however the Th1 cytokine changes observed were greater than the Th2 (51, 52). In accordance with this concept, Alizadeh *et al.*, (2006) reported an increase in IL-12 and IFN- γ production on splenocytes derived from animals exposed to BPA in an OVA-induced allergy model (53). Coupled with this effect, Menard *et al.*, (2014) evaluated the BPA effect on the specific immune response against the OVA antigen, they found that BPA increased the LTh percentages and the IFN- γ secretion (54). It is important to mention that the perinatal BPA exposure (day 9.5 of gestation until weaning), produced changes in cytokine expression, such as G-CSF, GM-CSF, IL-12p70, IL-1 α , IL-1 β , TNF- α and Rantes at serum levels (55). Moreover, this cytokine secretion pattern together with the IL-4 and IL-6 increase was also observed in splenocytes previously stimulated with Con-A. Also, the supernatants of cultured splenocytes were evaluated and the GM-CSF, IFN- γ and IL-17 expression was also increased after LPS stimulus. All the above confirms that BPA impacts on Th1 cytokines with a bias towards a Th17 type response (55).

In mice, BPA exposure administrated in their drinking water during the perinatal period, increased the expression of the transcription factor ROR γ t, and the cytokines IL-17, IL-21, IL-6 and IL-23 in a dose-dependent manner, in both males and females. (56). This is in agreement with previously reported data by Holladay *et al.*, 2010. Some studies have reported that BPA was able to generate T cell polarization, directing it to a Th2 response. With respect to this statement, in an *in vitro* model, lymphocytes exposed to BPA showed an induction of the GATA-3, IL-4 and IL-10 expression and a reduction of the T-box transcription factor (Tbet) expression, suggesting an induction of Th2 polarization (57). Similar effects were reported in activated T lymphocytes with respect to IL-4 expression (58).

Additionally, Miao *et al.*, (2008), using a gestational exposure scheme, reports that exposure to BPA generates a decrease in ER α expression in males, while the expression of this receptor is increased in females and the expression of Th1 cytokines (IL-2, IL-12, IFN- γ and TNF- α) was decreased in both males and females (59). Sawai *et al.* (2003) in *in vitro* experiments on splenocytes stimulated with Con-A, reported that males exposed to BPA, produced on average 40% less IFN- γ while females 28% less compared to controls (60). Regarding on regulatory T cells (Tregs) Ohshima *et al.*, (2007), reported that the perinatal administration of BPA generated a decrease in the Tregs number (61).

5.2. B lymphocytes and plasmatic cells

B lymphocytes (BL) are cells whose importance lies in humoral immunity, since they are able to differentiate into plasmatic cells (PCs), which are responsible for the production of antibodies such as IgA, IgG, IgE, IgD and IgM. In different studies, it has been observed that BPA may modify in a different way the production of antibodies by these cells. Yoshino *et al.*, (2003) using male and female DBA / 1J adult mice, report that BPA generates an increase in the proliferation of splenocytes, in addition to an increase in the production of antibodies in a dose-dependent manner (51). Similarly, in another study, it was reported that gestational exposure to BPA generated increased production of IgG1 and IgG2a antibodies, with predominance of IgG2a antibodies in male offspring (52).

Menard *et al.* (2014), reported that in female rats exposed perinatally to BPA there is an increase in anti-OVA IgG antibodies during adult life (54). Alizadeh *et al.* (2006), indicate that exposure to BPA, in female BALB/c mice, generates lower IgE antibody titers and higher levels of IgG2a (53). Similarly, Lee *et al.* (2003), indicate that there is an increase in IgE in female BALB/c mice exposed to BPA during adult life (58). In another study, using young mice of the strain NZB/NZW (systemic lupus erythematosus murine model), which were exposed to BPA, and later in adult life the splenocytes were

Table 1. Effect of BPA on susceptibility to different infections

Pathogen	Mode of administration	Dose	Effect	Reference
Influenza A	Perinatal	50 µg/kg/day	Temporary reduction of the degree of pulmonary inflammation	64
<i>Escherichia coli</i>	Prepuberal	5 mg/kg/day x 5 days	Increase the number of colonies forming units	65
<i>Leishmania major</i>	Adult	5.7, 11.4, 22.8 y 45.6 mg/kg one week before infection	Increase susceptibility in a dose-dependent manner	68
<i>Leishmania major</i>	Perinatal	1, 10 y 100 nM	Increase susceptibility in a dose-dependent manner	68
<i>Trichinella spiralis</i>	Adult	228 µg/mice	Decreased susceptibility	69
<i>Nippostrongylus brasiliensis</i>	Perinatal	5 µg/kg/day	Increase susceptibility	44
<i>Toxocara canis</i>	Perinatal	250 µg/kg/day	Increase susceptibility	72

obtained and stimulated with LPS, there is a decrease in the production of IgG2a in splenocyte culture supernatants (60). Yurino *et al.* (2004), using BWF1 mice (another model of systemic lupus erythematosus), which were ovariectomized at four weeks of age, and that were subsequently implanted with subcutaneous implants of BPA during four months, reported that the exposure to BPA generates an increase in the production of anti-RBC IgM autoantibodies by B1 cells, as well as ERα expression both *in vitro* and *in vivo* (62). Moreover, Goto *et al.* (2007), administering BPA in the drinking water in male and female mice with transgenic TCR and stimulated with OVA reported an increase in the production of IgG2a and IgA in supernatants of splenocyte cultures from these mice (63). Finally, Midoro-Horiuti *et al.* (2010), using an asthma model in BALB/c mice, indicated that perinatal exposure of BPA in the drinking water, increases serum levels of anti-OVA IgG (42).

6. EFFECTS OF BPA ON INFECTIONS

6.1. Virus

So far, there is only one report about the effects of BPA in the virus infection context. Roy *et al.* (2012), evaluated the perinatal BPA exposure on the immune response associated with the infection of influenza A virus during adult life on a murine model (Table 1). The report indicated that the perinatal exposure to BPA did not affect the specific adaptive

immune response against the influenza virus at the lung. However, this exposure to BPA temporarily reduced the lung inflammation associated to the infection- and also the expression of antiviral genes (IFN-γ and iNOS) in lung tissue: The study concluded that perinatal BPA exposure modulated the innate immune response in the adult mice, but not the adaptive response, which is fundamental for the elimination of the influenza virus (64).

6.2. Bacteria

In the context of bacterial infections, BPA exposure, affected the defense against *Escherichia coli* (*E. coli*) in young mice (Table 1). BPA decreased the ability of immune cells to eliminate *E. coli* at 24 hrs post-infection. In addition, BPA induced neutrophil migration to the peritoneal cavity and reduced their phagocytic capacity against the bacteria. Coupled with these effects, a reduction in macrophage and lymphocyte populations were also observed (65). Moreover, the analysis of bacteria number in the peritoneal cavity indicated that there was a greater number of Colony Forming Units (CFU) in animals exposed to BPA as compared with untreated group (65).

While not properly an infection, bacterial colonization of the gastrointestinal tract needs to keep a balance between symbiotic, commensal and pathogenic species and maintains a bidirectional relationship with the immune system. In recent years,

the microbiome has gained attention, since it has been demonstrated that it can affect not only the intestinal health, but also the metabolism and even neurological phenomena. In that regard, BPA exposure has also been shown to alter the gut microbiome. Lai *et al.* (2016), found a significant reduction in gut microbiota diversity in mice exposed to BPA through out the diet. Interestingly, this effect is similar to that induced by high fat and high sucrose diets. Furthermore, this altered microbiota was enriched in proteobacteria, which it is a known as a marker of dysbiosis (66). Moreover, another group reported that the BPA effect on microbiome reaches the second generation of bacteria, and that some bacteria -associated disease such as *Bacteroides*, *Mollicutes*, *Prevotellaceae*, *Erysipelotrichaceae*, *Akkermansia*, *Methanobrevibacter* and *Sutterella* increase in proportion in BPA treated animals (67).

6.3. Parasites

The BPA effect upon immune response against *Leishmania major* (*L. major*) infection has been reported. In this context, the prenatal (administration of BPA in the drinking water, two weeks before mating and one-week post-mating, or in the adult life one week before to infection with *L. major*) induced an increase in the inflammation in the food pad of mice in a dose-dependent manner in *L. major* infected mice (Table 1). The Treg cells number was also reduced at splenic level in both, mice exposed to BPA at prenatally or in the adult stage. Of note, animals exposed to BPA in the adult stage have an IL-4, IL-10 and IL-13 increased level expression after *L. major* infection. Similar effect was reported in animals exposed to BPA prenatally where an IL-4 and IFN- γ increase was found (68). BPA effect in *Trichinella spiralis* (*T. spiralis*) infection has also been reported. The model was performed in mice exposed to a single dose (228 μg per mice) of this compound after 22 hours of *T. spiralis* infection, then mice were sacrificed at 42 days post infection and the muscle larvae number were counted. The study reported that BPA decreased the larvae number, suggesting that BPA has a protective effect against to this nematode infection (69). Recently, we have demonstrated that neonatal (3 days of age) male and female syngeneic BALB/c mice exposed to a single dose of BPA, and that further were infected with the human nematode

Trichinella spiralis in the adulthood, harbour fewer parasitic loads on the duodenum. Protective effect of BPA was related to an immunomodulatory effect of BPA related to the specific immune response to the helminth (70). Similarly, Nava-Castro *et al.* (2020) demonstrated that neonatal treatment with a single dose of BPA, decreases parasites loads in the adulthood in BalB/c female mice intraperitoneally infected with the helminth parasite *Tania crassiceps*. Also, in this case, BPA had an specific immunomodulatory effect on the immune response (71). On the contrary, Ménard *et al.* (2014), using a perinatal administration model (day 15 of pregnancy until day 21 postnatal) of BPA in rats, administered in their drinking water and infecting afterwards with the parasite *Nippostrongylus brasiliensis* (*N. brasiliensis*), reported an increase in susceptibility to infection in young female offspring (25 days of age) that were exposed to BPA perinatally (44). Similarly, Del Río *et al.* (2020), report an increase in susceptibility to *Toxocara canis* (*T. canis*) infection in adult male rats exposed perinatally (day 5 of gestation at 21 days postnatal) at a dose of 250 μg / Kg / day of BPA. The increase in larvae number was related to an inadequate polarization of the immune response, where there was an increase in Th1 cytokines TNF- α and IFN- γ , while there was a decrease in Th2 cytokines IL-4 and IL-5, as well as specific anti-*T. canis* IgG antibodies (72).

As for the direct effects exerted by BPA on parasites, Tan *et al.* (2015), reported that this estrogenic compound increased mortality of *Caenorhabditis elegans* (*C. elegans*), also BPA accelerated its aging process by increasing mitochondrial and cytosolic oxidative stress, as well as ROS generation (73). Other report indicated that BPA exposure at concentrations of 1 to 10 μM on *C. elegans* embryos decreased the parasites' oviposition during adult life (74). On the other hand, Zhou *et al.* (2016), in a multigenerational study using *C. elegans* as a model, reported that BPA induced physiological changes over the four generations of parasites, depending on the EDC concentration exposure (0.001-10 μM). In the first generation, parasites had a lower growth, moved slowly and produced lower offspring than nematodes that were not exposed to BPA (75). This work also referred that long-term BPA exposure (10 days), generated

BPA contributes to the patho-physiology of infectious diseases

chronic toxicity affecting physiological indicators as body size, head contractions, body curvature and half-life; coupled with a greater stress response and a decrease in the population size (76).

7. CONCLUSIONS

Endocrine disrupting compounds modulate endogenous steroid responses and cell functions. Although most studies focus on their reproductive effects, their potential effects on immune cells and even more, on the immune response towards pathogens, should draw attention, given the expression of hormonal receptors by immune cells. Despite the fact that a lot of studies have evaluated the BPA effect with different variables such as BPA concentration, type of model, administered BPA dose, life stage or antigenic challenge used, one common element remains: BPA can differentially modulate the immune response. Sometimes, BPA treatment helps the host to defend it self against a pathogen, sometimes BPA treatment it is detrimental and helps the invader. Thus, BPA not always has to be considered an enemy, but a double edge sword, that depending on the context of the However, more studies are needed with the aim to elucidate the possible mechanisms by which this takes place.

8. ACKNOWLEDGMENTS

Financial support: Grant IN-209719 from Programa de Apoyo a Proyectos de Innovación Tecnológica (PAPIIT), Dirección General de Asuntos del Personal Académico (DGAPA), Universidad Nacional Autónoma de México (UNAM) and Grant FC2016-2125 from Fronteras en la Ciencia, Consejo Nacional de Ciencia y Tecnología (CONACYT), both to Jorge Morales-Montor. Grant IA 206220 to Víctor H del Río Araiza, from PAPIIT, DGAPA, UNAM. Grant IA 202919 to Karen Elizabeth Nava-Castro, also from PAPIIT, DGAPA, UNAM. Carmen T. Gómez de León is a recipient of a Post-Doctoral fellowship from Grant FC2016-2125, Fronteras en la Ciencia, Consejo Nacional de Ciencia y Tecnología (CONACYT).

9. REFERENCES

1. C Guzmán-Arriaga; E Zambrano. Endocrine disruptor compounds and their

role in the developmental programming of the reproductive axis. *Rev Investig Clin* 59, 73-81 (2007)

2. V Pergialiotis; P Kotrogianni; E Christopoulos-Timogiannakis; D Koutaki; G Daskalakis; N Papantoniou. Bisphenol A and adverse pregnancy outcomes: a systematic review of the literature. *J Matern Fetal Neonatal Med* 31, 3320-3327 (2018)
DOI: 10.1080/14767058.2017.1368076
PMid:28805116
3. T Sweeney. Is exposure to endocrine disrupting compounds during fetal/post-natal development affecting the reproductive potential of farm animals? *Domest Anim Endocrinol* 23, 203-209 (2002)
DOI: 10.1016/S0739-7240(02)00157-1
4. JJ Amaral Mendes. The endocrine disrupters: A major medical challenge. *Food Chem Toxicol* 40, 781-788 (2002)
DOI: 10.1016/S0278-6915(02)00018-2
5. W V. Welshons; SC Nagel; FS Vom Saal. Large effects from small exposures. III. Endocrine mechanisms mediating effects of bisphenol A at levels of human exposure. *Endocrinology* 147, 56-69 (2006)
DOI: 10.1210/en.2005-1159
PMid:16690810
6. GG Kuiper; JG Lemmen; B Carlsson; JC Corton; SH Safe; PT van der Saag; B van der Burg; J a. Gustafsson. Interaction of estrogenic chemicals and pytoestrogens with estrogen receptor beta. *Endocrinology* 139, 4252-4263 (1998)
DOI: 10.1210/endo.139.10.6216
PMid:9751507
7. EC Bonefeld-Jørgensen; M Long; M V.

- Hofmeister; AM Vinggaard. Endocrine-disrupting potential of Bisphenol A, Bisphenol A dimethacrylate, 4-n-nonylphenol, and 4-n-octylphenol *in vitro*: New data and a brief review. *Environ Health Perspect* 115, 69-76 (2007)
DOI: 10.1289/ehp.9368
PMid:18174953 PMCID:PMC2174402
8. K Pogrmic-Majkic; G Kosanin; D Samardzija Nenadov; S Fa; B Stanic; A Trninc Pjevic; N Andric. Rosiglitazone increases expression of steroidogenic acute regulatory protein and progesterone production through PPAR γ -EGFR-ERK1/2 in human cumulus granulosa cells. *Reprod Fertil Dev* 31, 1647-1656 (2019)
DOI: 10.1071/RD19108
PMid:31233701
 9. RM Sargis; DN Johnson; RA Choudhury; MJ Brady. Environmental endocrine disruptors promote adipogenesis in the 3T3-L1 cell line through glucocorticoid receptor activation. *Obesity (Silver Spring)* 18, 1283-8 (2010)
DOI: 10.1038/oby.2009.419
PMid:19927138 PMCID:PMC3957336
 10. K Moriyama; T Tagami; T Akamizu; T Usui; M Saijo; N Kanamoto; Y Hataya; A Shimatsu; H Kuzuya; K Nakao. Thyroid hormone action is disrupted by bisphenol A as an antagonist. *J Clin Endocrinol Metab* 87, 5185-90 (2002)
DOI: 10.1210/jc.2002-020209
PMid:12414890
 11. YB Wetherill; BT Akingbemi; J Kanno; JA McLachlan; A Nadal; C Sonnenschein; CS Watson; RT Zoeller; SM Belcher. *In vitro* molecular mechanisms of bisphenol A action. *Reprod Toxicol* 24, 178-198 (2007)
DOI: 10.1016/j.reprotox.2007.05.010
PMid:17628395
 12. P Alonso-Magdalena; AB Roper; S Soriano; M García-Arévalo; C Ripoll; E Fuentes; I Quesada; Á Nadal. Bisphenol-A acts as a potent estrogen via non-classical estrogen triggered pathways. *Mol Cell Endocrinol* 355, 201-207 (2012)
DOI: 10.1016/j.mce.2011.12.012
PMid:22227557
 13. JR Rochester. Bisphenol A and human health: A review of the literature. *Reprod Toxicol* 42, 132-155 (2013)
DOI: 10.1016/j.reprotox.2013.08.008
PMid:23994667
 14. S Lee; C Kim; H Shin; Y Kho; K Choi. Comparison of thyroid hormone disruption potentials by bisphenols A, S, F, and Z in embryo-larval zebrafish. *Chemosphere* 221, 115-123 (2019)
DOI: 10.1016/j.chemosphere.-2019.01.019
PMid:30639807
 15. MJ Kim; YJ Park. Bisphenols and Thyroid Hormone. *Endocrinol Metab (Seoul, Korea)* 34, 340-348 (2019)
DOI: 10.3803/EnM.2019.34.4.340
PMid:31884733 PMCID:PMC6935774
 16. R Rezg; S El-Fazaa; N Gharbi; B Mornagui. Bisphenol A and human chronic diseases: current evidences, possible mechanisms, and future perspectives. *Environ Int* 64, 83-90 (2014)
DOI: 10.1016/j.envint.2013.12.007
PMid:24382480
 17. DD Seachrist; KW Bonk; SM Ho; GS Prins; AM Soto; RA Keri. A review of the carcinogenic potential of bisphenol A, <http://www.ncbi.nlm.nih.gov/pubmed/264>

- 93093, (2016)
DOI: 10.1016/j.reprotox.2015.09.006
PMid:26493093 PMCID:PMC4783235
18. I Campesi; M Marino; A Montella; S Pais; F Franconi. Sex Differences in Estrogen Receptor α and β Levels and Activation Status in LPS-Stimulated Human Macrophages, (2017)
DOI: 10.1002/jcp.25425
PMid:27171902
19. Y Wang; L Wang; J Zhao; Z Qiao. Estrogen, but not testosterone, down-regulates cytokine production in nicotine-induced murine macrophage. *Methods Find Exp Clin Pharmacol* 27, 311-316 (2005)
DOI: 10.1358/mf.2005.27.5.893666
PMid:16082418
20. C-C Hong; M Shimomura-Shimizu; M Muroi; K Tanamoto. Effect of endocrine disrupting chemicals on lipopolysaccharide-induced tumor necrosis factor-alpha and nitric oxide production by mouse macrophages. *Biol Pharm Bull* 27, 1136-1139 (2004)
DOI: 10.1248/bpb.27.1136
PMid:15256756
21. U Yamashita; T Sugiura; Y Yoshida; E Kuroda. Effect of endocrine disrupters on macrophage functions *in vitro*. *J UOEH* 27, 1-10 (2005)
DOI: 10.7888/juoeh.27.1_1
PMid:15794588
22. X Lu; M Li; C Wu; C Zhou; J Zhang; Q Zhu; T Shen. Bisphenol A promotes macrophage proinflammatory subtype polarization via upregulation of IRF5 expression *in vitro*. *Toxicol In vitro* 60, 97-106 (2019)
DOI: 10.1016/j.tiv.2019.05.013
PMid:31108126
23. YH Kuan; FM Huang; YC Li; YC Chang. Proinflammatory activation of macrophages by bisphenol A-glycidyl-methacrylate involved NF κ B activation via PI3K/Akt pathway. *Food Chem Toxicol* 50, 4003-4009 (2012)
DOI: 10.1016/j.fct.2012.08.019
PMid:22939937
24. Y Liu; C Mei; H Liu; H Wang; G Zeng; J Lin; M Xu. Modulation of cytokine expression in human macrophages by endocrine-disrupting chemical Bisphenol-A. *Biochem Biophys Res Commun* 451, 592-598 (2014)
DOI: 10.1016/j.bbrc.2014.08.031
PMid:25128825
25. D Teixeira; C Marques; D Pestana; A Faria; S Norberto; C Calhau; R Monteiro. Effects of xenoestrogens in human M1 and M2 macrophage migration, cytokine release, and estrogen-related signaling pathways. *Environ Toxicol* 31, 1496-1509 (2016)
DOI: 10.1002/tox.22154
PMid:26011183
26. M Yang; W Qiu; B Chen; J Chen; S Liu; M Wu; KJ Wang. The *in vitro* immune modulatory effect of bisphenol a on fish macrophages via estrogen receptor ?? and Nuclear Factor-??B signaling. *Environ Sci Technol* 49, 1888-1895 (2015)
DOI: 10.1021/es505163v
PMid:25565130
27. JJ Segura; A Jiménez-Rubio; R Pulgar; N Olea; JM Guerrero; JR Calvo. *In vitro* Effect of the Resin Component Bisphenol A on Substrate Adherence Capacity of Macrophages. *J Endod* 25, 341-344 (1999)
DOI: 10.1016/S0099-2399(06)81168-4

28. J Young Kim; H Gwang Jeong. Down-regulation of inducible nitric oxide synthase and tumor necrosis factor- α expression by bisphenol A via nuclear factor- κ B inactivation in macrophages. *Cancer Lett* 196, 69-76 (2003)
DOI: 10.1016/S0304-3835(03)00219-2
PMid:24287822
29. JA Byun; Y Heo; YO Kim; MY Pyo. Bisphenol A-induced downregulation of murine macrophage activities *in vitro* and *ex vivo*. *Environ Toxicol Pharmacol* 19, 19-24 (2005)
DOI: 10.1016/j.etap.2004.02.006
PMid:21783458
30. T Ohnishi; T Yoshida; A Igarashi; M Muroi; KI Tanamoto. Effects of possible endocrine disruptors on MyD88-independent TLR4 signaling. *FEMS Immunol Med Microbiol* 52, 293-295 (2008)
DOI: 10.1111/j.1574-695X.2007.00355.x
PMid:18177342
31. J Yoshitake; K Kato; D Yoshioka; Y Sueishi; T Sawa; T Akaike; T Yoshimura. Suppression of NO production and 8-nitroguanosine formation by phenol-containing endocrine-disrupting chemicals in LPS-stimulated macrophages: involvement of estrogen receptor-dependent or -independent pathways. *Nitric oxide Biol Chem* 18, 223-8 (2008)
DOI: 10.1016/j.niox.2008.01.003
PMid:18252206
32. KH Kim; SM Yeon; HG Kim; HS Choi; H Kang; HD Park; TW Park; SP Pack; EH Lee; Y Byun; SE Choi; KS Lee; UH Ha; YW Jung. Diverse influences of androgen-disrupting chemicals on immune responses mounted by macrophages. *Inflammation* 37, 649-656 (2014)
DOI: 10.1007/s10753-013-9781-1
33. S Liu; C Pan; Y Tang; F Chen; M Yang; KJ Wang. Identification of novel long non-coding RNAs involved in bisphenol A induced immunotoxicity in fish primary macrophages. *Fish Shellfish Immunol* 100, 152-160 (2020)
DOI: 10.1016/j.fsi.2020.03.006
PMid:32147374
34. S Kovats. Estrogen receptors regulate innate immune cells and signaling pathways. *Cell Immunol* 294, 63-69 (2015)
DOI: 10.1016/j.cellimm.2015.01.018
PMid:25682174 PMCID:PMC4380804
35. H Guo; T Liu; Y Uemura; S Jiao; D Wang; Z Lin; Y Narita; M Suzuki; N Hirose; Y Ichihara; O Ishihara; H Kikuchi; Y Sakamoto; S Senju; Q Zhang; F Ling. Bisphenol A in combination with TNF- α selectively induces Th2 cell-promoting dendritic cells *in vitro* with an estrogen-like activity. *Cell Mol Immunol* 7, 227-34 (2010)
DOI: 10.1038/cmi.2010.14
PMid:20383177 PMCID:PMC4002911
36. L Pisapia; G Del Pozzo; P Barba; L Caputo; L Mita; E Viggiano; GL Russo; C Nicolucci; S Rossi; U Bencivenga; DG Mita; N Diano. Effects of some endocrine disruptors on cell cycle progression and murine dendritic cell differentiation. *Gen Comp Endocrinol* 178, 54-63 (2012)
DOI: 10.1016/j.ygcen.2012.04.005
PMid:22531466
37. U Švajger; MS Dolenc; M Jeras. *In vitro* impact of bisphenols BPA, BPF, BPAF and 17 β -estradiol (E2) on human monocyte-derived dendritic cell generation, maturation and function. *Int Immunopharmacol* 34, 146-154 (2016)

- DOI: 10.1016/j.intimp.2016.02.030
PMid:26945833
38. A Camarca; C Gianfrani; F Ariemma; I Cimmino; D Bruzzese; R Scerbo; S Picascia; V D'Esposito; F Beguinot; P Formisano; RV Valentino. Human peripheral blood mononuclear cell function and dendritic cell differentiation are affected by bisphenol-A exposure. *PLoS One* 11, 1-18 (2016)
DOI: 10.1371/journal.pone.0161122
PMid:27509021 PMCid:PMC4980038
39. M Chakhtoura; U Sriram; M Heayn; J Wonsidler; C Doyle; J-A Dinnall; S Gallucci; RA Roberts. Bisphenol A Does Not Mimic Estrogen in the Promotion of the *In vitro* Response of Murine Dendritic Cells to Toll-Like Receptor Ligands. *Mediators Inflamm* 2017, 2034348 (2017)
DOI: 10.1155/2017/2034348
PMid:28811679 PMCid:PMC5547709
40. H Watanabe; R Adachi; K Kusui; A Hirayama; T Kasahara; K Suzuki. Bisphenol A significantly enhances the neutrophilic differentiation of promyelocytic HL-60 cells. *Int Immunopharmacol* 3, 1601-1608 (2003)
DOI: 10.1016/S1567-5769(03)00182-6
41. T Reistad; E Mariussen; F Fonnum. The effect of a brominated flame retardant, tetrabromobisphenol-A, on free radical formation in human neutrophil granulocytes: The involvement of the MAP kinase pathway and protein kinase C. *Toxicol Sci* 83, 89-100 (2005)
DOI: 10.1093/toxsci/kfh298
PMid:15456914
42. T Midoro-Horiuti; R Tiwari; CS Watson; RM Goldblum. Maternal bisphenol a exposure promotes the development of experimental asthma in mouse pups. *Environ Health Perspect* 118, 273-277 (2010)
DOI: 10.1289/ehp.0901259
PMid:20123615 PMCid:PMC2831929
43. M He; T Ichinose; S Yoshida; H Takano; M Nishikawa; T Shibamoto; G Sun. Exposure to bisphenol A enhanced lung eosinophilia in adult male mice. *Allergy Asthma Clin Immunol* 12, 16 (2016)
DOI: 10.1186/s13223-016-0122-4
PMid:27087817 PMCid:PMC4832452
44. S Ménard; L Guzylack-Piriou; C Lencina; M Leveque; M Naturel; S Sekkal; C Harkat; E Gaultier; M Olier; R Garcia-Villar; V Theodorou; E Houdeau. Perinatal exposure to a low dose of bisphenol a impaired systemic cellular immune response and predisposes young rats to intestinal parasitic infection. *PLoS One* 9, 1-15 (2014)
DOI: 10.1371/journal.pone.0112752
PMid:25415191 PMCid:PMC4240706
45. E O'Brien; DC Dolinoy; P Mancuso. Bisphenol A at concentrations relevant to human exposure enhances histamine and cysteinyl leukotriene release from bone marrow-derived mast cells. *J Immunotoxicol* 11, 84-9 (2014)
DOI: 10.3109/1547691X.2013.800925
PMid:23782309 PMCid:PMC4030600
46. E O'Brien; DC Dolinoy; P Mancuso. Perinatal bisphenol A exposures increase production of pro-inflammatory mediators in bone marrow-derived mast cells of adult mice. *J Immunotoxicol* 11, 205-212 (2014)
DOI: 10.3109/1547691X.2013.822036
PMid:23914806 PMCid:PMC3983174
47. SL Klein; KL Flanagan. Sex differences in immune responses. *Nat Rev Immunol* advance on (2016)

- DOI: 10.1038/nri.2016.90
PMid:27546235
48. S Ansar Ahmed; WJ Penhale; N Talal. Sex hormones, immune responses, and autoimmune diseases. Mechanisms of sex hormone action. *Am J Pathol* 121, 531-51 (1985)
49. M Labib Salem; M Sohrab Hossain; K Nomoto. Mediation of the Immunomodulatory Effect of β -Estradiol on Inflammatory Responses by Inhibition of Recruitment and Activation of Inflammatory Cells and Their Gene Expression of TNF- α and IFN- γ . *Int Arch Allergy Immunol* 121:121 (2000)
50. J-Y Youn; H-Y Park; J-W Lee; I-O Jung; K-H Choi; K Kim; K-H Cho. Evaluation of the immune response following exposure of mice to bisphenol A: induction of Th1 cytokine and prolactin by BPA exposure in the mouse spleen cells. *Arch Pharm Res* 25, 946-53 (2002)
DOI: 10.1007/BF02977018
PMid:12510852
51. S Yoshino; K Yamaki; R Yanagisawa; H Takano; H Hayashi; Y Mori. Effects of bisphenol A on antigen-specific antibody production, proliferative responses of lymphoid cells, and TH1 and TH2 immune responses in mice. *Br J Pharmacol* 138, 1271-6 (2003)
DOI: 10.1038/sj.bjp.0705166
PMid:12711627 PMCID:PMC1573776
52. S Yoshino; K Yamaki; X Li; TAO Sai; H Takano; H Hayashi. Prenatal exposure to bisphenol A up-regulates immune responses, including T helper 1 and T helper 2 responses, in mice. *Immunology* 112, 495 (2004)
DOI: 10.1111/j.1365-2567.2004.01900.x
PMid:15196218 PMCID:PMC1782504
53. M Alizadeh; F Ota; K Hosoi; M Kato; T Sakai; M Satter. Altered allergic cytokine and antibody response in mice treated with Bisphenol A. *J Med Invest* 53, 70-80 (2006)
DOI: 10.2152/jmi.53.70
PMid:16537998
54. S Ménard; L Guzylack-Piriou; M Leveque; V Braniste; C Lencina; M Naturel; L Moussa; S Sekkal; C Harkat; E Gaultier; V Theodorou; E Houdeau. Food intolerance at adulthood after perinatal exposure to the endocrine disruptor bisphenol A. *FASEB J* 28, 4893-4900 (2014)
DOI: 10.1096/fj.14-255380
PMid:25085925
55. SD Holladay; S Xiao; H Diao; J Barber; T Nagy; X Ye; RM Gogal. Perinatal bisphenol A exposure in C57B6/129svj male mice: Potential altered cytokine/chemokine production in adulthood. *Int J Environ Res Public Health* 7, 2845-2852 (2010)
DOI: 10.3390/ijerph7072845
PMid:20717544 PMCID:PMC2922731
56. S Luo; Y Li; Y Li; Q Zhu; J Jiang; C Wu; T Shen. Gestational and lactational exposure to low-dose bisphenol A increases Th17 cells in mice offspring. *Environ Toxicol Pharmacol* 47, 149-158 (2016)
DOI: 10.1016/j.etap.2016.09.017
PMid:27693988
57. J Lee; K-T Lim. Plant-originated glycoprotein (36 kDa) suppresses interleukin-4 and -10 in bisphenol A-stimulated primary cultured mouse lymphocytes. *Drug Chem Toxicol* 33, 421-9 (2010)
DOI: 10.3109/01480541003739229
PMid:20553123

58. MH Lee; SW Chung; BY Kang; J Park; CH Lee; SY Hwang; TS Kim. Enhanced interleukin-4 production in CD4+ T cells and elevated immunoglobulin E levels in antigen-primed mice by bisphenol A and nonylphenol, endocrine disruptors: Involvement of nuclear factor-AT and CA2+. *Immunology* 109, 76-86 (2003)
DOI: 10.1046/j.1365-2567.2003.01631.x
PMid:12709020 PMCID:PMC1782943
59. S Miao; Z Gao; Z Kou; G Xu; C Su; N Liu. Influence of bisphenol a on developing rat estrogen receptors and some cytokines in rats: a two-generational study. *J Toxicol Environ Health A* 71, 1000-1008 (2008)
DOI: 10.1080/15287390801907467
PMid:18569609
60. C Sawai; K Anderson; D Walser-Kuntz. Effect of bisphenol A on murine immune function: Modulation of interferon-gamma, IgG2a, and disease symptoms in NZB X NZW F1 mice. *Environ Health Perspect* 111, 1883-1887 (2003)
DOI: 10.1289/ehp.6359
PMid:14644661 PMCID:PMC1241761
61. Y Ohshima; A Yamada; S Tokuriki; M Yasutomi; N Omata; M Mayumi. Transmaternal exposure to bisphenol a modulates the development of oral tolerance. *Pediatr Res* 62, 60-64 (2007)
DOI: 10.1203/PDR.0b013e3180674dae
PMid:17515845
62. H Yurino; S Ishikawa; T Sato; K Akadegawa; T Ito; S Ueha; H Inadera; K Matsushima. Endocrine disruptors (environmental estrogens) enhance autoantibody production by B1 cells. *Toxicol Sci* 81, 139-147 (2004)
DOI: 10.1093/toxsci/kfh179
PMid:15166399
63. M Goto; Y Takano-Ishikawa; H Ono; M Yoshida; K Yamaki; H Shinmoto. Orally administered bisphenol A disturbed antigen specific immunoresponses in the naïve condition. *Biosci Biotechnol Biochem* 71, 2136-43 (2007)
DOI: 10.1271/bbb.70004
PMid:17827700
64. A Roy; SM Bauer; BP Lawrence. Developmental exposure to bisphenol a modulates innate but not adaptive immune responses to influenza a virus infection. *PLoS One* 7, 1-12 (2012)
DOI: 10.1371/journal.pone.0038448
PMid:22675563 PMCID:PMC3366985
65. Y Sugita-Konishi; S Shimura; T Nishikawa; F Sunaga; H Naito; Y Suzuki. Effect of Bisphenol A on non-specific immunodefenses against non-pathogenic Escherichia coli. *Toxicol Lett* 136, 217-227 (2003)
DOI: 10.1016/S0378-4274(02)00388-0
66. K-P Lai; Y-T Chung; R Li; H-T Wan; CK-C Wong. Bisphenol A alters gut microbiome: Comparative metagenomics analysis. *Environ Pollut* 218, 923-930 (2016)
DOI: 10.1016/j.envpol.2016.08.039
PMid:27554980
67. AB Javurek; WG Spollen; SA Johnson; NJ Bivens; KH Bromert; SA Givan; CS Rosenfeld. Effects of exposure to bisphenol A and ethinyl estradiol on the gut microbiota of parents and their offspring in a rodent model. *Gut Microbes* 7, 471-485 (2016)
DOI: 10.1080/19490976.2016.1234657
PMid:27624382 PMCID:PMC5103659
68. H Yan; M Takamoto; K Sugane. Exposure to bisphenol A prenatally or in adulthood promotes TH2 cytokine

- production associated with reduction of CD4+CD25+ regulatory T cells. *Environ Health Perspect* 116, 514-519 (2008)
DOI: 10.1289/ehp.10829
PMid:18414636 PMCID:PMC2290985
69. X Tian; M Takamoto; K Sugane. Bisphenol A promotes IL-4 production by Th2 cells. *Int Arch Allergy Immunol* 132, 240-247 (2003)
DOI: 10.1159/000074305
PMid:14646385
70. KE Nava-Castro; H Solleiro-Villavicencio; VH del Río-Araiza; M Segovia-Mendoza; A Pérez-Torres; J Morales-Montor. Sex-associated protective effect of early bisphenol-A exposure during enteric infection with *Trichinella spiralis* in mice. *PLoS One* 14, 1-14 (2019)
DOI: 10.1371/journal.pone.0218198
PMid:31291264 PMCID:PMC6619665
71. KE Nava-Castro; C Togno-Peirce; MI Palacios-Arreola; VH Del Río-Araiza; M Segovia-Mendoza; R Hernández-Bello; J Morales Montor. Exposure to the endocrine disruptor compound Bisphenol A induces protection through modulation of the immune response against the helminth parasite *Taenia crassiceps*. *Par Immunol* 42, e12733 (2020)
72. VH Del Río-Araiza; MI Palacios-Arreola; KE Nava-Castro; NY Pérez-Sánchez; R Ruíz-Manzano; M Segovia-Mendoza; MI Girón-Pérez; MS Navidad-Murrieta; J Morales-Montor. Perinatal exposure to bisphenol A increases in the adulthood of the offspring the susceptibility to the human parasite *Toxocara canis*. *Environ Res* 184, 1-10 (2020)
DOI: 10.1016/j.envres.2020.109381
PMid:32199324
73. L Tan; S Wang; Y Wang; M He; D Liu. Bisphenol A exposure accelerated the aging process in the nematode *Caenorhabditis elegans*. *Toxicol Lett* 235, 75-83 (2015)
DOI: 10.1016/j.toxlet.2015.03.010
PMid:25819108
74. MD Mersha; BM Patel; D Patel; BN Richardson; HS Dhillon. Effects of BPA and BPS exposure limited to early embryogenesis persist to impair non-associative learning in adults. *Behav Brain Funct* 11, 27 (2015)
DOI: 10.1186/s12993-015-0071-y
PMid:26376977 PMCID:PMC4573949
75. D Zhou; J Yang; H Li; Q Lu; Y Di Liu; KF Lin. Ecotoxicity of bisphenol A to *Caenorhabditis elegans* by multigenerational exposure and variations of stress response *in vivo* across generations. *Environ Pollut* 208, 767-773 (2016)
DOI: 10.1016/j.envpol.2015.10.057
PMid:26561446
76. D Zhou; J Yang; H Li; C Cui; Y Yu; Y Liu; K Lin. The chronic toxicity of bisphenol A to *Caenorhabditis elegans* after long-term exposure at environmentally relevant concentrations. *Chemosphere* 154, 546-551 (2016)
DOI: 10.1016/j.chemosphere.2016.04.011
PMid:27085314

Key Words: Review, Environmental Pollution, Environmental Medicine, Endocrine Disruptors, Bisphenol A, Immunoregulation, Infection, Review

Send correspondence to: Jorge Morales-Montor, Departamento de Inmunología, Instituto de Investigaciones Biomedicas, Universidad Nacional Autonoma de Mexico, AP 70228, CP 04510, Ciudad de Mexico, Mexico, Tel: 52555-56223158, Fax: 525556223369, E-mail: jmontor66@biomedicas.unam.mx